TX 551

### U. S. DEPARTMENT OF AGRICULTURE.

OFFICE OF EXPERIMENT STATIONS—BULLETIN NO. 116.

A. C. TRUE, Director.

# DIETARY STUDIES IN NEW YORK CITY

### 1896 and 1897.

#### W. O. ATWATER, Ph. D.,

Professor of Chemistry, We legan University; Chief of Nutrition Investigations. Office of Experiment Stations,

AND

#### A. P. BRYANT, M. S.,

Assistant in Nutrition Investigations.



LIBRARY OF COMO BIVISION OF BOOM

WASHINGTON:

GOVERNMENT PRINTING OFFICE.


1902.






Class TX551
Book A92









### U. S. DEPARTMENT OF AGRICULTURE.

OFFICE OF EXPERIMENT STATIONS—BULLETIN NO. 116.

A. C. TRUE, Director.

5-20

# DIETARY STUDIES IN NEW YORK CITY

IN

### 1896 and 1897.

BY

#### W. O. ATWATER, Ph. D.,

Professor of Chemistry, Wesleyan University; Chief of Nutrition Investigations, Office of Experiment Stations,

AND

#### A. P. BRYANT, M. S.,

Assistant in Nutrition Investigations.



#### WASHINGTON:

GOVERNMENT PRINTING OFFICE.

1902.

000032

Monograph

TX551

#### OFFICE OF EXPERIMENT STATIONS.

A. C. TRUE, Ph. D., Director.

E. W. Allen, Ph. D., Assistant Director and Editor of Experiment Station Record.

C. F. Langworthy, Ph. D., Editor and Expert on Foods and Animal Production.

#### NUTRITION INVESTIGATIONS, MIDDLETOWN, CONN.

W. O. Atwater, Ph. D., Chief of Nutrition Investigations.

C. D. Woods, B. S., Special Agent at Orono, Me.

F. G. BENEDICT, Ph. D., Physiological Chemist.

R. D. Milner, Ph. B., Editorial Assistant.

7. m. S. Oct. 5. de,

#### LETTER OF TRANSMITTAL

U. S. DEPARTMENT OF AGRICULTURE,
OFFICE OF EXPERIMENT STATIONS,

Washington, D. C., July 2, 1902.

SIR: In order to secure satisfactory dietary standards, it is necessary to know the amounts of food actually consumed by a considerable number of persons of different food habits and living under different circumstances as regards occupation and environment. The studies made under the auspices of this Department have furnished much information regarding the dietaries of farmers, professional men, laborers, mechanics, college and university students (both men and women), persons of limited incomes living in large cities, etc. The studies reported herewith were made with families for the most part in poor circumstances, living in the thickly congested districts of New York City. The statistics regarding the families studied and their food consumption were gathered by Dr. Isabelle Delaney. Almost without exception the families were of the type frequently receiving help from charitable organizations. In some cases the income was as large as many families live upon in comparative comfort. In other cases the income was very small.

The data recorded show that the lack of proper food was frequently due to ignorance in buying and preparing it, or to some similar cause. Before the condition of families like many of those studied can be permanently improved, it is necessary to learn the errors which are commonly made in their domestic economy. Studies like those reported are a help in this regard. Indeed, the results already obtained have been made use of by the New York Society for the Improvement of the Condition of the Poor, which cooperated with this Department in carrying on this investigation.

The investigations were made under the immediate direction of the chief of the nutrition investigations, Prof. W. O. Atwater, of Wesleyan University.

The report is submitted with the recommendation that it be published as Bulletin No. 116 of this Office.

A. C. TRUE,


Director.

Hon. James Wilson, Secretary of Agriculture.



## CONTENTS.

	Page.
Introduction	7
Dietary studies in New York City	8
Details of the studies here reported	9
Dietary study of a gripman's family (No. 154)	10
Dietary study of a longshoreman's family (No. 155)	12
Dietary study of a plumber's family (No. 158)	14
Dietary study of a washerwoman's family (No. 159)	16
Dietary study of a truckman's family (No. 160)	17
Dietary study of a washerwoman's family (No. 161)	19
Dietary study of a longshoreman's family (No. 162)	21
Dietary study of a carpenter's family (No. 166)	22
Dietary study of a carpenter's family (No. 167)	24
Dietary study of a housekeeper's family (No. 168)	25
Dietary study of a caretaker's family (No. 170)	27
Dietary study of a tanner's family (No. 171)	29
Dietary study of a foundryman's family (No. 172)	30
Dietary study of a truckman's family (No. 177)	():)
Dietary study of a longshoreman's family (No. 178)	34
Dietary study of a carpenter's family (No. 180)	36
Dietary study of a sail rigger's family (No. 183)	38
Dietary study of a day laborer's family (No. 185)	39
Dietary study of a fruit yender's family (No. 186)	40
Dietary study of a watchman's family (No. 187)	42
Dietary study of a bookbinder's family (No. 188)	44
Dietary study of a butcher's family (No. 192)	45
Dietary study of a sail rigger's family (No. 193)	46
Dietary study of a washerwoman's family (No. 194)	48
Dietary study of a stableman's family (No. 195)	49
Dietary study of a truckman's family (No. 196)	51
Dietary study of a huckster's family (No. 197)	52
Dietary study of a longshoreman's family (No. 198)	54
Dietary study of a carpenter's family (No. 199)	56
Dietary study of a painter's family (No. 200)	57
Dietary study of an expressman's family (No. 201)	59
Dietary study of a waiter's family (No. 204)	61
Dietary study of a landlord's family (No. 205)	
Dietary study of a caretaker's family (No. 206)	
Dietary study of a sailor's family (No. 209)	
Dietary study of a housekeeper's family (No. 210)	
Summary and discussion.	
Pecuniary economy of food purchased	
Summary of amounts of nutrients and energy per man per day	
Conclusion.	
Amendix	Mar. 1



## DIETARY STUDIES IN NEW YORK CITY IN 1896 AND 1897.

#### INTRODUCTION.

The most reliable data concerning the food consumption of people of different nationality, age, sex, and occupation, living under different financial and hygienic conditions, are obtained by means of dietary These have been carried on quite actively in the United States during recent years, to some extent by independent investigators, but more extensively by individuals and institutions in different parts of the country working in cooperation with the United States Department of Agriculture. A number of these investigations were made with families with very limited incomes, more particularly those living in the congested districts of some of the larger cities, so that considerable information concerning the normal and usual food consumption of such persons has accumulated. The information thus obtained is of much importance, for besides being of direct value to those interested in improving the conditions of the less favored classes of the community, it forms an indispensable part of the general data of an adequate and comprehensive science of nutrition, especially in establishing dietary standards.

The present bulletin reports thirty-six studies made under the auspices of the Department of Agriculture among people with very limited means living in the more congested districts of New York City. They are a continuation of previous studies, which were similar in character to studies carried on about the same time in Pittsburg and in Chicago. Studies made among Mexican families in New Mexico and among negroes in Alabama and Virginia show the food consumption of people with very limited means but not crowded together in cities.

Among other studies made among families of very limited means in

a U. S. Dept. Agr., Office of Experiment Stations Bul, 46.

<sup>&</sup>lt;sup>b</sup> U. S. Dept, Agr., Office of Experiment Stations Bul. 52.

c U. S. Dept. Agr., Office of Experiment Stations Bul. 55.

d U. S. Dept. Agr., Office of Experiment Stations Buls. 40 and 54.

e U. S. Dept. Agr., Office of Experiment Stations Bul. 38.

f U. S. Dept. Agr., Office of Experiment Stations Bul. 71,

this country that might be especially mentioned here is an extensive series carried on by Miss Amelia Shapleigh, with the aid and supervision of Mrs. Ellen H. Richards, among poor families in Philadelphia and Chicago," but not yet published in detail. Some interesting studies were also undertaken in Hartford, Conn., by Miss Helen M. Hall under the joint auspices of the Hartford School of Sociology and the Storrs (Conn.) Experiment Station.<sup>b</sup>

Investigations of a similar nature have been carried on in other countries. Among these the recent dietary studies among laboring classes in Edinburgh by Drs. Noël, Paton, J. Craufurd Dunlop, and Elsie M. Inglis, and those of laborers families in York, England, made by Mr. B. Seebohm Rowntree, are of particular interest in this connection, both because of their large intrinsic importance and because they were carried on by the same methods as those of the studies here reported and among families in much the same circumstances.

#### DIETARY STUDIES IN NEW YORK CITY.

As already suggested, the particular purpose of the investigations reported on the following pages and in the bulletin previously mentioned e was to obtain reliable data concerning the food consumption of the classes of people living in the crowded districts of New York City. It seemed most advisable to make dietary studies among selected families that were believed to be representative of the regions in which they lived. To make such a selection, however, and to secure accurate and reliable statistics, it was necessary that the work be done by some one who was in sympathy with the people and familiar with their daily life. Those who are brought officially and personally into direct contact with them and have the opportunity and the means for studying their modes of life it is believed can best collate the facts regarding their food, what they buy, how much they pay for it, how they cook and eat it, and how in any or all of these respects improvements can be made. Such favorable conditions were secured in these investigations by the cooperation of the New York Association for the Improvement of the Condition of the Poor, one of the oldest and largest benevolent associations in the United States, which had the advantages of large resources, long experience, and close connection with the people in the congested quarters of the city. The selection of the families to be studied and the collection of the various statistics regarding their circumstances and their food consumption devolved upon Dr. Isabelle Delaney, whose long experience in mission work, and

<sup>&</sup>quot;" A study of dietaries." Partial report of Dutton Fellow, College Settlements Association, 1892–93.

<sup>&</sup>lt;sup>b</sup>Conn. (Storrs) Sta. Rpt. 1896, p. 117.

<sup>&</sup>lt;sup>c</sup>Diet of Laboring Classes in Edinburgh.

d Poverty, a Study of Town Life, p. 222.
 e U. S. Dept. Agr., Office of Experiment Stations Bul. 46.

especially as the family physician of a very large number of people in the regions referred to, gave her unusual opportunities for understanding the people and their conditions, experiences, and ideas, while her sympathy with them and their confidence in her secured the freest admission to their homes to herself and anyone she brought with her.

In addition to those already reported (see p. 7), studies of thirty-six families were made during 1896 and 1897, the details of which are here given. The families selected represented many nationalities and occupations. The range in total income per family was from an amount not sufficient to buy the actual necessities of life to an amount equal to that upon which families in other communities have been found to live comfortably. In some instances the persons studied were slovenly and thriftless, taking little interest in their homes. Other families, though ignorant, were willing and anxious to learn how to improve their habits of living.

#### DETAILS OF THE STUDIES HERE REPORTED.

The studies were carried on, and the final results were calculated, according to methods described in detail in previous bulletins.<sup>a</sup> The data sought included (1) the nationality, age, sex, and occupation of the different members of the family and their general physical condition; (2) the income of the family; (3) the expenditures for rent and for food; (4) the kind, quality, and quantity of the food consumed; and (5) the number of meals taken by each person present during the study. From these data and the standard tables showing the composition and fuel value of the different food materials used the quantities of nutrients and energy consumed per man per day were computed. It was then possible, upon comparison with the results of similar studies elsewhere and with recognized standards, to judge whether the families studied were properly nourished, and whether they were wise in their selection and purchase of food; also to point out, in many instances, how a more nutritious diet might have been obtained at the same cost, or one equally nutritious for less.

The results of the thirty-six dietary studies are given in the following pages. The text and the tables for each study contain all the data from which the cost and quantities of nutrients and energy per man per day have been computed. The final figures represent very nearly, if not exactly, the amounts actually consumed, as, although the waste was not determined, in the majority of cases it was observed to be very small, if indeed there was any at all. The circumstances of most of the families positively prohibited their throwing away any edible material.

No analyses were made in connection with these studies. The composition of each of the different food materials used was assumed to

<sup>&</sup>lt;sup>a</sup> U. S. Dept. Agr., Office of Experiment Stations Bul. 46, and others given in list on cover of this bulletin.

be the same as that given in tables of average con.position of food materials." The percentages of nutrients and the fuel values employed for calculating the amounts of nutritive ingredients in these dietaries are given in Table 47 in the Appendix. The numbers in the column headed "Reference number" in this table are the same as those given in parentheses in connection with the weights and cost of the food materials in the table for each dietary study, and thus serve to indicate the data used to calculate the quantities of nutrients in the different materials.

The fuel values of the nutrients were calculated by the use of the same factors as were employed in previous bulletins. Somewhat smaller factors have been proposed recently, but the older factors have been here retained in order that these studies may be directly comparable with those previously reported.

In several instances in the following pages there is given in the discussion of the dietary a table showing the price per pound of the various food materials purchased, the amounts of protein and energy in 1 pound, and the total quantity of each material and of protein and energy that could be obtained for 10 cents at the given price per pound, as well as the total amount expended for each food during the period of study. By the figures in such a table it is possible to form an estimate of the pecuniary economy of the different materials at the prices paid.

#### DIETARY STUDY OF A GRIPMAN'S FAMILY (NO. 154).

This study was made with a family consisting of the father, the mother, and one child. The father was a strong, healthy man, American born, weighing 230 pounds, and was employed as gripman on a cable car. He was on duty from noon until midnight, and usually took one meal each day away from home. The mother, Italian born, was strong and healthy, and weighed 175 pounds. She was a capable housewife. The boy, 14 years of age, weighed 75 pounds, and was rather delicate. The income of the family was \$12 per week, of which \$3.50 was taken by the father to pay for his meals away from home. Thirteen dollars a month rent was paid for three rooms, two well-lighted and one on an air shaft. The family was well dressed and carried considerable insurance, but saved nothing otherwise.

The study began June 1, 1896, and continued ten days. The number of meals taken was as follows:

	Meals.
Man	22
Woman (30 meals × 0.8 meal of man), equivalent to	24
Boy, 14 years (30 meals $\times$ 0.8 meal of man), equivalent to	24
Total number of meals taken equivalent to	70
Equivalent to one man twenty-three days.	

a U. S. Dept. Agr., Office of Experiment Stations Bul. 28.

b By the present writers. See Conn. (Storrs) Sta. Rpt. 1899, p. 110.

Table 1.— Weights and cost of food and nutrients in dietary study No. 154.

Kinds, amounts, and cost of food for ten days, a	Cost, nutrients, and fuel value of food per day.				
which and cost of foot for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.  Beef: Sirloin, 2.38 pounds, 42 cents (33); bologna,	Cents.	Grams.	Grams.	Grams.	Calories.
0.25 pound, 5 cents (1); gelatin, 0.44 pound, 7 cents (15). Veal: Chops, 4.93 pounds, 60 cents (54)	5.0	36			324
5.25 pounds, 63 cents (66).  Poultry: Chicken, 3.37 pounds, 40 cents (83).  Fish, etc.: Cod, fresh, 1.69 pounds, 25 cents (90); her-	2.9	16 13	37 11		410 155
rings, smoked, 1 pound, 10 cents (96); oysters, 2.19 pounds, 20 cents (101) Eggs, 6.34 pounds, 83 cents (115) Butter, 2.75 pounds, 47 cents (118) . Cheese, 1.75 pounds, 25 cents (120)	2. 4 3. 6 2. 0 1. 1	12 18 1 9	2 13 46 12	1	72 194 432 153
Milk, 24.02 pounds, 62 cents (124)	2.7	16	159	24	2,080
VEGETABLE FOOD.	==	121	100		2,000
Cereals: Rice, 0.50 pound, 4 cents (130); bread, 14.50 pounds, 64 cents (134); cake, 2.94 pounds, 28 cents (142); macaroni, 2.50 pounds, 15 cents (158).  Sugar, 6 pounds, 32 cents (169)  Vegetables: Asparagus, 1 pound, 10 cents (174); lettuce, 4.36 pounds, 22 cents (193); onions, 2.51	4.8 1.4	37	10	233 118	1, 200 484
pounds, 6 cents (1.95); potatoes, 14.20 pounds, 45 cents (204); radishes, 3.24 pounds, 13 cents (208); tomatoes, canned, 13.01 pounds, 51 cents (216) Fruits: Bananas, 0.69 pound, 5 cents (225); cherries,	6.4	12	1	73	357
0.87 pound, 10 cents (227); strawberriès, 2.76 pounds, 25 cents (239)	1.8	1	1	10	54
Total vegetable food	14.4	50 [	12	434	2,095
Total food	35, 8	171	171	460	4, 175

 $<sup>^{\</sup>circ}$  The numbers in parentheses after each food material in this and succeeding tables refer to corresponding numbers in Table 47, p. 79.

The diet in this study was unusually large; but, on the other hand, the father and mother were unusually heavy and the man was at rather hard muscular work. Based on the standard of 125 grams of protein and 2,500 calories of energy for a man at moderate work, it would appear that the family had more food than they needed. Taking into account, however, their weight and the occupation of the father, it is probable that they were receiving about what would satisfy the physiological demands of the body.

According to the statistics of the study the food cost about \$5.75 per week, or, approximately, half the income. This sum was equivalent to 36 cents per man per day, which, though perhaps not excessive considering the variety of food materials and the quantity of nutrients and energy obtained, was nevertheless larger than was necessary. Yet there was evidently careful management, as is shown by the variety in the food obtained at the price paid and also by the fact that there was no waste, "left-over" pieces being carefully utilized.

The relative economy of the food materials purchased by this family is illustrated by the figures in the following table:

Table 2.—Cost of food materials per pound, and amounts of protein and energy obtained for 10 cents in each, in dietary study No. 154.

	In 1 pound. Amount bought for 10 cents.					Total	
Kind of food material.	Price per pound.	Protein.	Fuel value.	Total food.	Protein.	Fuel value.	amount ex- pended during study,
Beef, sirloin Veal chops Smoked ham Chicken Fish, fresh (assumed as cod) Herring, smoked Oysters Eggs. Butter Cheese Milk Bread Cake Macaroni Sugar Green vegetables: Asparagus, lettuce, onions, radishes Potatoes	Cents. 17. 6 12. 2 12. 0 11. 9 14. 8 10. 0 9. 1 13. 1 17. 1 14. 3 2. 6 4. 4 9. 5 6. 0 5. 3 4. 6 3. 2	Pound. 0.165 199 142 193 165 205 060 148 010 259 033 092 063 134	Calories. 825 825 1, 675 1, 045 325 750 230 720 3, 605 1, 950 325 1, 215 1, 675 1, 665 1, 860	Pounds. 0.57 82 83 84 68 1.00 1.10 76 6.59 70 8.85 2.27 1.055 1.67 1.89	Pound. 0.09 16 12 16 11 21 07 11 .18 .18 .21 .07 .22 .63 .07	Cullories. 556 675 1,395 890 215 755 566 24,105 1,250 2,745 1,760 2,780 3,485	Cents. 42 60 63 40 25 10 20 83 47 25 62 64 28 15 32
Tomatoes, canned	9.3	. 012	105	2.55	.03	265 280	51 40

The prices paid for various food materials were, as a rule, very reasonable. Fresh bread at 4.4 cents a pound was cheap, and, together with macaroni at 6 cents a pound, formed by far the most economical source of both protein and energy in the diet. As compared with the protein and energy obtained for the money expended for bread and macaroni, it is interesting to note the amounts obtained in 10 cents' worth of green vegetables, canned tomatoes, and fruits. During the time of the study \$1.42 was expended for these latter materials, the amount of nutrients obtained being about the same as in 15 cents' worth of bread. The meats used were the more expensive cuts. Had they bought the cheaper cuts of meat and used less oysters and fewer eggs, the cost of the diet might have been reduced materially. If, in addition, some of the money expended for canned tomatoes, fresh fruit, and green vegetables had been used to purchase more economical food, the diet might have been still further reduced in cost, and at the same time have been equally or more nutritious.

#### DIETARY STUDY OF A LONGSHOREMAN'S FAMILY (NO. 155).

The family in this study was believed to be typical of a large class living "from hand to mouth," buying food in small quantities as wanted for each meal. The members of the family were the father, the mother, the grandmother, and four boys, aged, respectively, 13, 11, 8, and 3 years. The father, Irish born, was a longshoreman, weighing 160 pounds, and rather quiet in disposition and stolid. His income varied with the amount of work he could get. During the period of study he was earning about \$8 per week. The mother, weighing 130 pounds,

was thrifty and hard working. She took care of the halls in the building in which the family lived. For this service she was allowed rooms which would probably have rented for about \$13 per month; she also did washing and cleaning when the father was out of work and it was necessary for her to earn money. The grandmother was strong and well and did considerable housework. The boys were small for their age and sickly, and appeared to be insufficiently nourished. The oldest was employed as errand boy and earned \$1.50 a week. The family occupied four very small, dark rooms, in only one of which was a window that would admit much light or air, the other three opening upon an air shaft. The dimensions of each of the two bedrooms were 6 by 7 feet.

The study began June 2, 1896, and continued ten days. The number of meals taken was as follows:

	Mears.
Man	30
Two women (60 meals $\times$ 0.8 meal of man), equivalent to	. 48
Two boys, 13 and 11 years (60 meals $\times$ 0.64 meal of man), equiva	-
lent to	. 36
One boy, 8 years (30 meals $\times$ 0.5 meal of man), equivalent to	. 15
One boy, 3 years (30 meals $\times$ 0.4 meal of man), equivalent to	. 12
Visitor	. 2
Total number of meals taken equivalent to:	143
Equivalent to one man forty-eight days.	

Table 3.—Weights and cost of food and nutrients in dietary study No. 155.

Winds amounts and seat of fined for ten dura	Cost, nutrients, and fuel value of food per day.				per man
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.  Beef: Steak, chuck, 5 pounds, 48 cents (27); soup, fore shank, 1.51 pounds, 9 cents (23); corned bris-	Cents,	Grams.	Grams.	Grams.	Calories.
ket, 9.5 pounds, 60 cents (4); tripe, 5.88 pounds, 30 cents (38); suet, 0.38 pound (37). Pork: Ham, boiled, 0.88 pound, 21 cents (64); spare-	3.1	33	31		424
rib, roast, 5.62 pounds, 53 cents (81); pig's head, 2.06 pounds, 10 cents (70); lard, 0.50 pound, 6 cents (69). Fish: Cod, fresh, 6.5 pounds, 28 cents (89). Eggs, 1 pound, 10 cents (116). Butter, 3.69 pounds, 73 cents (118). Milk, 23.62 pounds, 56 cents (124).	1.8 .6 .2 1.5 1.2	13 7 1 0 7	30 0 1 30 9	11	332 29 13 279 158
Total animal food	8.4	61	101	11	1,235
VEGETABLE FOOD.			*		Name of the last o
Cereals: Bread, 7.5 pounds, 39 cents (134); bread, stale, 17 pounds, 26 cents (138). Sugar: 6.5 pounds, 39 cents (169). Vegetables: Beans, string, 2 pounds, 7 cents (177); cabbage, 4.82 pounds, 10 cents (179); cucumbers,	1.4	24	3	124 62	634 254
0.87 pound, 5 cents (186); onions, 1.19 pounds, 3 cents (195); potatoes, 31.18 pounds, 55 cents (204); spinach, 2.5 pounds, 8 cents (214)	1.8	9	1	60	292
Total vegetable food	4.0	33	4	246	1,180
Total food	12.4	94	105	257	2, 415

a This factor was used instead of the more common one because the boys were small for their ages.

The expenditures for food in this dietary were on the whole well made. In spite of the fact that food was purchased in small amounts, and that there was an unnecessary though pleasing variety of vegetables, the cost per man per day was but 12.4 cents. The ration was, however, deficient in protein and especially lacking in energy. Had half the 33 cents expended for fresh vegetables, other than potatoes, been expended for dried beans, and the 39 cents used to purchase fresh bread been spent for "stale" bread (that is, bread a day old but not so old as to be unpalatable or unwholesome) at the price paid for the latter, the ration per man per day would have been increased by 25 grams of protein and 510 calories of energy. While this would have reduced the variety in the diet to some extent it would have perhaps improved the general condition of the children, who seemed insufficiently nourished.

#### DIETARY STUDY OF A PLUMBER'S FAMILY (NO. 158).

This family consisted of the father, an American, 28 years old, weighing 140 pounds; the mother, 26 years old, weighing 125 pounds; and two daughters, respectively 4 and 2 years old and rather small for their age. The grandfather, aged 74 years, weighing 150 pounds; a great uncle, 75 years old, weighing 160 pounds; and an uncle, 28 years, weighing 135 pounds, also lived with them. The father, a steamfitter's helper, was a strong, healthy man, but was idle at the time and did not seem anxious to work. His wife was thrifty and neat and a good manager. She and the two children appeared to be poorly nourished. The grandfather was in good health but without ambition. The great uncle earned his pocket money but contributed nothing to his support. The uncle, who was a plumber, paid \$5 a week for support of himself and the great uncle. He took his dinners away from home. The family occupied three rear rooms, for which they paid \$8.50 a month rent. One room was lighted from the rear yard, the other two opened on a hall and an air shaft. Two beds and a lounge served as sleeping quarters for the seven people. They lived in the easiest manner possible, set no table, bought their food by the meal, cooked it in the simplest manner, sat out of doors until late at night, and slept late in the morning. The food purchased was of poor quality, the milk being especially so.

The study began July 21, 1896, and continued ten days. The number of meals taken was as follows:

Men (four)	Meals.
Woman (29 meals $\times$ 0.8 meal of man), equivalent to	. 23
Two children, 4 and 2 years (60 meals × 0.4 meal of man), equived lent to.	
Total number of meals taken equivalent to	158

Equivalent to one man fifty-three days.

Table 4.—Weights and cost of food and nutrients in dietary study No. 158.

	Cost, ni	utrients, ar	nd fuel va per day.		per man
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy drates.	Fuel value.
ANIMAL FOOD.  Beef: Brisket, corned, 6.31 pounds, 39 cents (4); steak, chuck, 1.50 pounds, 15 cents (27); steak, round, 2 pounds, 22 cents (29); tripe, 1 pound, 6 cents (39). Mutton: Shoulder, 1.75 pounds, 8 cents	Cents.	Grams.	Grams.	Grams.	Calories.
(50); breast, 0.80 pound, 8 cents (40). Pork: Chops, 0.80 pound, 10 cents (60); shoulder, smoked, 5.25 pounds, 43 cents (79); bacon, 2 pounds, 22 cents (59); lard, 0.13 pound, 2 cents (69). Fish: Clams, 1 pound, 10 cents (85).	. 2	8 1	26	1	291 275 8 45
Eggs, 3.34 pounds, 40 cents (114) Butter, 2.75 pounds, 36 cents (118) Milk, 16.92 pounds, 47 cents (124).	1.0	5	20 6	7	186 105
Total animal food.	6.2	39	77	8	910
VEGETABLE FOOD.  Cereals: Bread, 20.56 pounds, 54 cents (138); cake, 1 pound, 10 cents (142).  Sugar, 7.75 pounds, 46 cents (169).  Vegetables: Cabbage, 3.69 pounds, 9 cents (179); corn, 2.25 pounds, 8 cents (184); onions, 1.07 pounds, 4 cents (195); potatoes, 19.45 pounds, 32 cents (204);	1.2	20	3	100 66	520 270
tomatoes, fresh, 5 pounds, 10 cents (215); tomatoes, canned, 2 pounds, 7 cents (216)	1.3	5	1	39	190
Total vegetable food	3, 4	25	4	205	980
Total food	9.6	64	si si	213	1,890

The amounts of protein and energy per man per day in this dietary were but little more than half of what is called for by the ordinary standard for a man at moderate work. Although the father was out of work at the time, and therefore required less food than if he had been actively employed, and two of the other men in the family did no work, still if the results given represent the average food consumption of the family it is not surprising that some of them appeared to be undernourished.

Considerable improvement in the nutritive value of the diet could have been made by diminishing the amounts of some of the foods selected and the substitution of others not used at all. For instance, animal foods were purchased in larger amounts and greater variety than was necessary, over three-fifths of the total expenditure having been for such materials, while but about one-eighth was for the cereals, which constitute the most economical source of nutriment ordinarily obtainable. They used no legumes, and no oatmeal, wheat, or other cereal foods, except bread and cake. The cabbage, corn, onions, and tomatoes cost more than the potatoes used, though they furnished less than half as much protein, and but little more than quarter the energy obtained in the potatoes. Had half of the \$2.15 expended for meat, fish, and eggs been expended for bread, corn meal, oatmeal, dried beans and peas, and the like, the quantity of nutrients in the diet

would have been greatly increased, while the cost would have remained the same. With proper cooking, the diet thus modified would doubtless have been no less attractive than the usual fare.

#### DIETARY STUDY OF A WASHERWOMAN'S FAMILY (NO. 159).

The members of the family here studied were the mother, 38 years of age, and six children—four girls, aged respectively 17, 14, 5, and 3 years, and two boys, aged respectively 10 and 7 years. The weights of all except the two younger girls were respectively 130, 140, 125, 85, and 65 pounds. The mother, English born, was refined and intelligent, her first husband having been a clergyman; her second marriage, however, had been unfortunate. The members of the family were not strong and found it difficult to obtain sufficient food for proper nourishment. The mother endeavored to support the family by washing and by house cleaning. A daughter, not living at home, contributed \$5 a month toward the rent. A sailor, 19 years old, weighing 150 pounds, boarded with the family during three days of the study, paying \$2.15. Provisions were bought for cash by the meal at small markets. There was no visible waste and but very little refuse. The family occupied three well-lighted rooms for which they paid \$10 a month. The rent was low for the locality, owing to the fact that the building was notorious for crimes that had been committed in it, and the rooms were not in demand.

The study began July 21, 1896, and continued ten days. Three of the children went to the country before the close of the study. The number of meals taken was as follows:

	Meals.
Woman (30 meals $\times$ 0.8 meal of man), equivalent to	
Two girls, 17 and 14 years old (19 meals $\times$ 0.7 meal of man), equi-	va-
lent to	13
Boy, 10 years old (30 meals $\times$ 0.6 meal of man), equivalent to	18
Boy, 7 years old (28 meals $\times$ 0.5 meal of man), equivalent to	14
Two girls, 5 and 3 years old (30 meals $\times$ 0.4 meal of man), equi	13.7
lent to	12
Boarder	7
Visitor	2
Total number of meals taken equivalent to	90

Equivalent to one man thirty days.

Table 5.—Weights and cost of food and nutrients in dietary study No. 159.

Kinds, amounts, and cost of food for ten days.		Cost, nutrients, and fuel value of food per day.				
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy-drates.	Fuel value.	
ANIMAL FOOD.	Cents.	Canana	Charana	Grams,	Chalonian	
Beef: Liver, 2 pounds, 14 cents (18); sausage, bolo- gna, 0.25 pound, 5 cents (1); shank, fore, 3 pounds, 20 cents (23); steak, round, 2 pounds, 24 cents (29); steak, skirt, 2 pounds, 14 cents (35); steak, sirioin,	Cemes.	tifums,	Grams,	Grams,	caiories.	
1 pound, 12 cents (33) Pork: Bacon, 1 pound, 12 cents (59); chops, 2,5	3.0	26	17		270	
pounds, 20 cents (61)	1.1	6	19		200	
salmon, 2 pounds, 40 cents (104) Eggs, 0.32 pound, 5 cents (114) Butter, 1 pound, 20 cents (118)	- 1.8 2 6	11	4 1 13		86 13 120	
Milk, 11.83 pounds, 36 cents (124)	1.2	6 3	7 3	9 20	12 12	
Total animal food	8, 6	53	61	29	93	
VEGETABLE FOOD,						
Cereals: Flour, 1 pound, 5 cents (131); oatmeal, 3.51 pounds, 8 cents (128); bread, rye, 6 pounds, 17 cents (136); bread, wheat, 17.15 pounds, 36 cents (138); soda biscuit, 2.5 pounds, 25 cents (139); cakes, 1 pound, 10 cents (142); crackers, 6 pounds, 10 cents (142); crackers, 10 cent						
26 cents (153); apple pie, 0.25 pound, 10 cents (160). sugar, 4 pounds, 22 cents (169). Vegetables: Canned corn, 2 pounds, 9 cents (185); onions, 1 pound, 4 cents (195); potatoes, 7.9 pounds,	4.6	60	26	329 61	1,83 25	
15 cents (204); radishes, 0.25 pound, 2 cents (209); tomatoes, 3 pounds, 6 cents (215); canned tomatoes, 4 pounds, 12 cents (216)	1.6	5	1	33	16	
Fuit: Apples, 2 pounds, 7 cents (221); currants, 2 pounds, 7 cents (228); pears, 2 pounds, 10 cents (236).	.8	1		. 11	5	
Total vegetable food	7.7	66	27	434	2,30	
Total food	16.3	119	91	463	3,23	

The results of this study may be taken as a typical illustration of the fact that a varied and nutritious diet can be obtained at a comparatively small cost. The quantities of protein and energy per man per day in the ration were not greatly below the standard, and, considering the cost of the food, 16.3 cents, perhaps the ration could not be much improved upon. There was a considerable variety of animal food, the greatest expenditure for any one item being for canned salmon, which, it is interesting to note, furnished less nutriment than was obtained in the fore shank of beef for half as much money. Cereal foods were purchased in considerable variety and quantity. There was also a variety in the vegetables and fruits, though no sort was used in large amounts. By reducing the variety of vegetables and fruits the cost of the ration could have been still further reduced without materially diminishing its nutritive value. If sufficient care were taken in cooking, the less expensive diet could be made about as attractive as that costing more.

#### DIETARY STUDY OF A TRUCKMAN'S FAMILY (NO. 160).

This study was carried on in a family somewhat above the average in intelligence and thrift for the vicinity. It included the father, Amer-

ican born, of Scotch-Irish descent, 29 years old; the mother, 24 years old, and three children—a boy of 5, a girl of 3, and an infant of 1½ years of age, all of whom were in good health. They weighed 165, 138, 45, and 35 pounds, respectively, the weight of the infant not being ascertained. The father, a truckman, earned \$14 a week. The family dressed neatly, and made a good appearance. They occupied two large, well-lighted rooms, for which \$10 a month rent was paid. The rooms were furnished very simply, but were kept very neat and clean. Food was purchased for cash at the large or small markets, according to the nature of the material. Ice was used, and food was well cared for. There was practically no waste. The woman was thrifty, and though not an expert cook, was bright, and eager to learn how to improve the character of the diet. She had already profited much from lessons in domestic management given by one of the prominent local charitable organizations.

The study began October 20, 1896, and continued ten days. The number of meals taken was as follows:

	means,
Man	. 26
Woman (30 meals $\times$ 0.8 meal of man), equivalent to	. 24
Two children, 5 and 3 years old (46 meals $\times$ 0.4 meal of man), equiv	-
alent to	. 18 .
Infant, 1½ years old, equivalent to	. 9
Total number of meals taken equivalent to	_ 77
Equivalent to one man twenty-six days.	

Table 6.— Weights and cost of food and nutrients in dietary study No. 160.

	Cost, n	alue of foo	food per man		
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.	Cents.	Grams.	Grams.	Grams.	Calories.
Beef: Round steak, 3 pounds, 48 cents (29); fore shank, 4 pounds, 28 cents (23); plate, corned, 3.25 pounds, 20 cents (7). Mutton: Leg, 4.26 pounds, 96 cents (43); chops, 1 pound, 14 cents (44).  Pork: Sparerib, 2.06 pounds, 17 cents (81).  Fish: Mackerel, salt, 1.63 pounds, 17 cents (99); sal-	7.9	42 6	49 11		628 127
mon, canned, 0.50 pound, 18 cents (105) mont, canned, 0.50 pound, 18 cents (105) Butter, 3.13 pounds, 52 cents (118) Milk, 6 pounds, 12 cents (124) Milk, condensed, 3.25 pounds, 23 cents (125)	.5	7 1 3 5	6 46 4 5	5 31	84 432 70 194
Total animal food	13.3	64	121	36	1,535
VEGETABLE FOOD.  Cereals: Oatmeal, 1.75 pounds, 8 cents (129); macaroni, 1 pound, 10 cents (158); bread, 10 pounds, 50 cents (134); biscuit, 2.25 pounds, 10 cents (139); buns, 7 pounds, 29 cents (140); cake, coffee, 2 pounds, 20 cents (143).  Sugar, 2.5 pounds, 15 cents (169).  Vegetables: Cabbage, 8 pounds, 12 cents (179); onions, 0.56 pound, 2 cents (195); peas, dry, 2 pounds, 6 cents (200); potatoes, 10.69 pounds, 24 cents (204); corn, canned, 1 pound, 10 cents (185); peas, canned, 1 pound, 10 cents (199); tomatoes,	4.9	39	21	234 44	1, 315 181
canned, 2 pounds, 9 cents (216).  Fruit: Jelly, currant, 1 pound, 10 cents (232)	2.8	17	2 1	71 12	380 59
Total vegetable food	8.6	56	24	361	1,935
Total food	21.9	120	145	397	3,470

The results of this dietary study indicate that the family were obtaining about the normal quantity of protein and energy in their daily food. The cost of the ration—22 cents per man per day—was not excessive, although it might easily have been reduced. The expenditure of 96 cents for a leg of lamb secured but little more than half the protein and much less than half the fat that was obtained for a similar amount expended for cheap cuts of beef. Cereal products were used in considerable variety. The price of the bread-5 cents per poundwas higher than that paid by a number of the families studied. If food had been bought in larger quantities, and a portion of the money expended for meat had been used to increase the quantity of cereals. the nutritive value of the ration would have been increased with little or no diminution of its variety and palatability. The variety of vegetables might perhaps also have been reduced and only the more economical kinds purchased without decreasing appreciably the palatability of the ration.

#### DIETARY STUDY OF A WASHERWOMAN'S FAMILY (NO. 161).

This study was carried on with the same family as in study No. 159, after the return of the children from an outing in the country. The income during the study was \$5.

The study began August 6, 1896, and continued ten days. The number of meals taken was as follows:

	Meats.
Woman (30 meals $\times$ 0.8 meal of man), equivalent to	. 24
Girl, 17 years old (30 meals $\times$ 0.7 meal of man), equivalent to	21
Boy, 10 years old (30 meals $\times$ 0.6 meal of man), equivalent to	18
Boy, 7 years old (30 meals $\times$ 0.5 meal of man), equivalent to	15
Two girls, 5 and 3 years old (60 meals $\times$ 0.4 meal of man), equiva-	a-
lent to	. 24
Total number of meals taken equivalent to	102

Table 7.— Weights and cost of food and nutrients in dietary study No. 161.

Equivalent to one man thirty-four days.

	Cost, nutrients, and fuel value of food per mar per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy-drates.	Fuel value.	
ANIMAL FOOD.  Beef: Flank, 1.50 pounds, 12 cents (35); fore shank, 2.50 pounds, 14 cents (23); corned, canned, 1 pound,	Cents.	Grams.	Grams.	Grams.	Calories.	
20 cents (6). Lamb: Chops, 1.31 pounds, 18 cents (41); chops, mutton, 0.81 pound, 15 cents (45)	2, 3	16	17		22	
Pork: Chops, 2.24 pounds, 25 cents (60); salt, 4 pounds, 28 cents (72). Fish: Salmon, canned, 0.50 pound, 10 cents (105);	1.6	6	55		53	
sardines, 1.50 pounds, 9 cents (107)		6	3		5	
Eggs, 0.13 pound, 2 cents (117)	$\binom{(a)}{1.5}$	(a)	(0)		(a) 29	
Milk, 18.70 pounds, 51 cents (124)	1.5	8	10	13	17	
Wilk, condensed, 4.25 pounds, 31 cents (125)		5	5	30	19	
Total animal food	8.4	42	121	43	1,47	

Table 7.—Weights and cost of food and nutrients in dietary study No. 161—Continued.

	Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.	
VEGETABLE FOOD.  Cereals: Oatmeal, 3.50 pounds, 8 cents (128); bread, 20 pounds, 52 cents (138); bread, rye, 4 pounds, 12	Cents.	Grams.	Grams,	Grams.	Calories.	
cents (136); biscuit, soda, 6 pounds, 20 cents (139); crackers, soda, 3 pounds, 14 cents (156); cakes, mixed, 1 pound, 5 cents (142); pie, apple, 0.25 pound, 5 cents (160)	3.4	51	23			
Sugar, 6 pounds, 33 cents (169). Vegetables: Beans, string, 1 pound, 5 cents (178); cabbage, 3 pounds, 5 cents (179); corn, canned, 1 pound, 5 cents (185); onions, 0.50 pound, 2 cents	1.0			80	328	
(195); potatoes, 11.23 pounds, 22 cents (204); tomatoes, 2.50 pounds, 5 cents (215) Fruits: Bananas, 0.94 pound, 7 cents (226); currants, fresh, 1 pound, 3 cents (228); muskmelon, 0.50	1.4	5	1	35	173	
pound, 3 cents (234)	.3			4	15	
Total vegetable food	6.1	59	24	403	2, 115	
Total food	14.5	101	145	416	3,590	

This study shows, as did the earlier one with the same family, unusually careful management of the household expenses. The variety of the food was considerable, and the quantity of nutrients obtained per man per day as large as has been found in many studies of farmers, mechanics, and other workingmen in the United States, though the cost was less than 15 cents per day. This is an instance of what can be done by careful management.

The following table, showing the amounts of protein and energy purchased for 10 cents in some of the more important food materials used by this family, illustrates the relative economy of their purchases:

Table 8.—Cost of food materials per pound, and amounts of nutrients and energy obtained for 10 cents in each, in dietary study No. 161.

	In 1 pound. Amounts bought for 10 cents.				r 10 cents.	Total amount	
Kind of food material.	Price per pound.	Protein.	Fuel value.	Total food.	Protein.	Fuel value.	ex- pended during study.
Beef: Flank Fore shank Corned, canned Lamb chops		Pound. 0.170 .128 .263 .187 .160	Calories. 1, 115 545 1, 280 1, 540 1, 695	Pounds. 1, 25 1, 79 , 50 , 73 , 54	Pound. 0.21 .22 .14 .14 .09	Calories. 1,460 975 560 1,120 915	Cents. 12 14 20 18 15
Pork: Chops, edible portion. Salt. Salmon, canned. Sardines. Butter. Milk. Milk, condensed. Oatmeal	11. 2 7. 0 20. 0 6. 0 18. 6 2. 7 7. 3	. 166 . 019 . 218 . 237 . 010 . 033 . 088 . 161	1,580 3,670 915 950 3,605 325 1,520 1,860	. 90 1. 43 . 50 1. 67 . 54 3. 70 1. 37 4. 35	. 15 . 03 . 11 . 39 . 12 . 12 . 70	1, 410 5, 250 450 1, 580 1, 935 1, 180 2, 090 8, 135	25 28 10 9 51 51 31 8
Bread: Wheat. Rye. Biseuits, soda. Crackers, soda Cakes, mixed Pie, apple Sugar.	3.0 3.3 4.7 5.0 20.0	. 109 . 090 . 093 . 098 . 005 . 005	1, 215 1, 180 1, 730 1, 925 1, 675 1, 270 1, 860	3, 85 3, 33 3, 00 2, 14 2, 00 , 50 1, 82	. 42 . 30 . 28 . 21 . 13 . 02	4, 820 3, 940 5, 185 4, 130 3, 365 640 3, 380	52 12 20 14 5 5 33
Green vegetables: String beans, cabbage, onions, and tomatoes Potatoes. Fruit: Bananas, currants, muskmelons.	2.0	. 022		4, 17 5, 00 1, 88	.06	575 1, 975	17 22 13

The canned corned beef at 20 cents a pound was very high priced, and the lamb chops at 18½ cents, though more economical than the corned beef, were also high. The most economical food material was outmeal, which was purchased at 2.3 cents per pound, while stale bread at 2.6 cents per pound was also very economical and was purchased in considerable amounts. The amount spent for green vegetables and fruit was perhaps no larger than health demanded. It is interesting to compare the quantities of protein and energy obtained for 10 cents by this family with corresponding amounts in dietary study No. 154 (p. 10).

#### DIETARY STUDY OF A LONGSHOREMAN'S FAMILY (NO. 162).

The family consisted of the husband, a German, 33 years old, weighing 155 pounds; the mother, a Scotch woman, 35 years old, weighing 130 pounds; two sons of the latter, one 19 years old, weighing 125 pounds, the other 12 years of age, and a woman boarder 18 years old. The weights of the younger son and the boarder were not ascertained. The income of the family was larger than most of those studied. The father earned from \$18 to \$25 a week at his work as longshoreman, and the mother about \$5 a week selling papers. The older son was apprenticed to a printer and earned \$3 a week. The boarder was out of work at the time of the study and was seeking employment, meanwhile her board remained in arrears. The family occupied three rooms, paying \$8.50 a month rent. They dressed well and had a considerable sum of money laid by. About the only form of recreation in the warm weather was an occasional day at some seaside resort. Food was purchased in small quantities, although ice was used continually.

The study began August 6, 1896, and continued ten days. The number of meals taken was as follows:

Λ	1eals
Two men	- 60
Two women (60 meals $\times$ 0.8 meal of man), equivalent to	48
Boy, 12 years old (30 meals $\times$ 0.6 meal of man), equivalent to	18
Total number of meals taken equivalent to	126

Equivalent to one man forty-two days.

Table 9.— Weights and cost of food and nutrients in dietary study No. 162.

Kinds, amounts, and cost of food for ten days.	Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of 1000 for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.	
ANIMAL FOOD.  Beef: Round steak, 6 pounds, 70 cents (29); sirloin steak, 1.50 pounds, 23 cents (33). Veal: Breast, 4.12	Cents.	Grams,	Grams.	Grams.	Calories.	
pounds, 35 cents (53)	3.0	23	17		253	
Pork: Ham, smokèd, 5.13 pounds, 64 cents (66); bacon, 0.38 pound, 5 cents (59). Fish: Bluefish, fresh, 3.51 pounds, 25 cents (84); cod-	1.7	.8	21		228	
fish, fresh, 3.51 pounds, 24 cents (89); sturgeon, 37 pounds, 10 cents (111)	1.4	9			37	
Eggs, 5.73 pounds, 82 cents (114)	1.9	8	7		98	
Butter, 4 pounds, 75 cents (118) Cheese, 1.38 pounds, 31 cents (120)	1.8	1 5	37 6		348 76	
Cheese, limburger, 0.37 pound, 10 cents (123)	( <i>u</i> )	(a)	(a)	(a)	(et)	
Milk, 8.39 pounds, 23 cents (121)		3 3	3	17 17	70 110	
Total animal food	11.9	60	95	22	1,220	
VEGETABLE FOOD.						
Cereals: Bread, 25.87 pounds, 97 cents (134); cake, coffee, 1.50 pounds, 10 cents (143) Sugar, 4 pounds, 18 cents (169)	2. 6 . 4	27	5	159 43	809 176	
Vegetables: Cabbage, 10 pounds, 18 cents (179); potatoes, 24 pounds, 40 cents (204); tomatoes, 6.50 pounds, 14 cents (215); turnips, 2.37 pounds, 5 cents (218).	1, 8	8		59	275	
Total vegetable food	4.8	35	5	261	1,260	
Total food	16.7	95	100	283	2,480	

a Amounts too small to affect results.

The quantity of nutrients and energy per man per day obtained by this family seems small in view of the fairly severe work of the father. The income was large enough to warrant a more liberal diet, and there was no apparent reason why the family should not be well nourished, so it may be that the diet was sufficient for their needs. The cost was moderate, especially considering the relatively large proportion of protein as compared with the energy. Had the amount expended for cereal food been doubled and expended as judiciously as in the previous dietary study, the quantity of protein per man per day could have been increased to about 135 grams and the energy to 3,650 calories, although the total cost would have been but 19.3 cents.

#### DIETARY STUDY OF A CARPENTER'S FAMILY (NO. 166).

This study was with the same family as that in dietary study No. 31, the details of which have been published in a former report." They were at this time in rather better circumstances than at the time the first study was made. The members of the family were all German born, and comprised the father 47 and the mother 37 years of age, and three children—a boy of 18, a girl of 14, and a boy of 11 years. All

were apparently in excellent health, and weighed, respectively, 157, 192, 150, 143, and 83 pounds. The father was a carpenter and earned \$10 a week when he had steady work. The mother acted as house-keeper or janitor in the building in which they lived and received in payment for her services the rent of four rooms, worth about \$12 a month. The older son was employed by an electrician and earned \$7 a week; he spent 15 cents a day for his lunch. The daughter earned \$3 a week as salesgirl; the younger boy went to school. The food purchased was of good quality and there was no avoidable waste.

The study began September 3, 1896, and continued ten days. The number of meals taken was as follows:

	Meals.
Two men <sup>a</sup>	_ 55
Woman (30 meals $\times$ 0.8 meal of man), equivalent to	. 24
Girl, 14 years old (30 meals $\times$ 0.7 meal of man), equivalent to	. 21
Boy, 11 years old (30 meals $\times$ 0.6 meal of man), equivalent to	. 18
Total number of meals taken equivalent to	. 118
Fauivalent to one man thirty-nine days	

Table 10.—Weights and cost of food and nutrients in dietary study No. 166.

	Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein,	Fat.	Carbohy- drates.	Fuel value.	
ANIMAL FOOD.  Beef: Steak, chuck, 1.75 pounds, 15 cents (27); steak, round, 9.50 pounds, 95 cents (29); steak, round, chopped, 1.25 pounds, 20 cents (31); steak, sirloin, 0.75 pound, 15 cents (33); fore shank, 4 pounds, 23	Cents.	Grams,	Grams.	Grams.	Calories.	
cents (23); liver sausage, 0.50 pound, 10 cents (19).	4.6	36	23		. 361	
Pork: Loin, 3.75 pounds, 38 cents (61); ham, 0.75 pound, 15 cents (66); lard, 1.50 pounds, 12 cents (69). Eggs, 4.94 pounds, 75 cents (114). Butter, 3 pounds, 62 cents (118). Milk, 53.25 pounds, 81.36 (124).	1.9	7 9		31	317 92 279 441	
Total animal food			115			
VEGETABLE FOOD.  Cereals: Flour, 11.50 pounds, 37 cents (131); bread, 14.25 pounds, 57 cents (134); rolls, Vienna, 1 pound, 5 cents (165); doughnuts, 6 pounds, 30 cents (157).  Sugar, 7 pounds, 35 cents (169).  Vegetables: Beans, 2 pounds, 10 cents (175); cabbage, 5 pounds, 7 cents (179); corn, 1.50 pounds, 5 cents, (184); onions, 1 pound, 3 cents (195); potatoes, 35.69 pounds, 45 cents (201); sulad, 3 pounds, 7 cents (211); sauerkraut, 2 pounds, 12 cents (212); soup greens, 10 t rounds, 15 cents (195); tentatoes, 6 counds.	3.3	36	18	232	1, 267 332	
1.94 pounds, 15 cents (189); tomatoes, 6 pounds, 11 cents (215).  Fruits: Apples, 3.25 pounds, 8 cents (222)	3.0	18	i)	104 4	519 17	
Total vegetable food	7,5	54	20	421	2, 135	
Total food	20.7	126	135	452	3,625	

<sup>&</sup>quot;As the meal taken away from home by the young man was only a lunch, it was assumed that he was absent from home for only 5 meals, rather than 10, and would eat at the other meals at home sufficient to make up the difference.

The results obtained in this study indicate that the family was receiving ample nourishment, perhaps a little more than was absolutely necessary; the cost, however, was not excessive. In the previous study of this same family they consumed 148 grams of protein and 3,825 calories of energy per man per day, quantities considerably in excess of those here found. The cost of the diet in the previous study was 23 cents per man per day as compared with 20.7 in this. In the discussion of the earlier study it was pointed out that a reduction in the food might easily have been made, and that such a change would probably not be a disadvantage. A still further reduction in cost might have been made in the present dietary, while still keeping its nutritive value equally high, by a wiser selection of vegetables. One of the highest priced of the articles of animal food purchased was sirloin steak, but the 15 cents thus expended furnished very much more nutriment than was obtained for the same sum expended for soup greens.

#### DIETARY STUDY OF A CARPENTER'S FAMILY (NO. 167).

The family consisted of the father, German born, 47 years of age; the mother, also German, 52 years of age; and four sons, aged respectively 20, 19, 15, and 7 years. The weights of the members of the family were respectively 220, 180, 120, 110, 75, and 47 pounds. The father was a carpenter and usually had steady work. At the time of the study he was recovering from an illness, and received \$6 a week from a benefit organization. The oldest son was a janitor in a clubhouse, the second son was a glassworker; neither earned large wages, vet each paid \$4 a week board. These two boys got their lunches each working day away from home. Since these were only light meals, it has been assumed that each one was present at 25 full meals during the study. The third son was learning the printer's trade; he paid \$2.75 a week toward his support. The mother was well trained in household management. The table was neat and inviting, the food well prepared, and there was little or no waste. The rent of the four well-lighted rooms occupied was \$6 per month. The rooms were comfortably furnished, and the family dressed better than was to be expected from their income.

The study began September 2, 1896, and continued ten days. The number of meals taken was as follows:

N N	eals.
Three men	80
Woman (30 meals × 0.8 meal of man), equivalent to	24
Boy, 15 years old (30 meals × 0.8 meal of man), equivalent to	
Boy, 7 years old (30 meals × 0.5 meal of man), equivalent to	
Total number of meals taken equivalent to	143

Equivalent to one man forty-eight days.

Table 11.—Weights and cost of food and nutrients in dietary study No. 167.

Kinds, amounts, and cost of food for ten days.	Cost, nutrients, and fuel value of food per man per day.					
All the state of t		Protein.	Fat.	Carbohy- drates.	Fuel value.	
ANIMAL FOOD,	Cents.	Grams.	Grams,	Grams.	Calories.	
Beef: Round steak, 3 pounds, 40 cents (29); steak, chopped, 3 pounds, 30 cents (31); loin, 2.50 pounds, 20 cents (34); shoulder, corned, edible portion, 4.50 pounds, 40 cents (5); suet, 0.12 pound (37). Veal: Leg, 4 pounds, 36 cents (56).	3.5	31	28		400	
Pork: Shoulder, smoked, 3.25 pounds, 23 cents (79); salt pork, 1 pound, 6 cents (72) Fish: Cod, dry, 0.88 pound, 7 cents (91); mackerel,	. 6	4	16		165	
fresh, 3 pounds, 16 cents (98).  Eggs, 5.65 pounds, 70 c nts (114).  Butter, 1.50 pounds, 33 cents (118).	1.5	5 7	1 6 12		30 85 110	
Milk, 6.12 pounds, 15 cents (124). Milk, condensed, 3 pounds, 27 cents (125)	.3	3	2 2	3 15	40 95	
Total animal food	7.6	55	67	18	925	
VEGETABLE FOOD.					7.7.7	
Cereals: Flour, 1.50 pounds, 5 cents (131); farina, 1 pound, 4 cents (127); bread, 4.50 pounds, 19 cents (134); bread, rye, 5.44 pounds, 25 cents (136); sugar buns, 7.13 pounds, 35 cents (140); sweet buns, 7.75 pounds, 35 cents (141); rolls, 0.75 pound, 5 cents (166). Sugar, 5 pounds, 29 cents (149)	2,7	23	10	145 47	782 193	
Vegetables: Cabbage, 4.13 pounds, 5 cents (179); corn, green, 0.50 pound, 5 cents (184); greens, 0.19 pound, 1 cent (188); onious, 1.19 pounds, 3 cents (195); potatoes, 40.49 pounds, 51 cents (204); tomatoes, 17						
pounds, 22 cents (215) Fruit: Bananas, 0.17 pound, 2 cents (225); grapes, 3.50	1.8	11	1	81	385	
pounds, 10 cents (230)	. 2			5	20	
Total vegetable food	5, 3	34	11	278	1,380	
Total food	12.9	89	78	296	2,305	

The cost of food per man per day in this study was very small, amounting to but 13 cents. On the other hand, the ration was scanty, even taking into account the fact that the father was recovering from sickness at the time and doubtless ate considerably less food than when at active work. One of the most expensive purchases as regards the nutritive return was tomatoes. Twenty-two cents expended for this vegetable furnished about three-quarters of the protein and energy obtained for 5 cents in wheat flour. Nevertheless, evidence of careful management is marked throughout the study. Meats were purchased in large amounts and considerable variety, but the cost was moderate. It must be remembered, however, that, generally speaking, the meats are a much more expensive source of nourishment than the cereals. The variety and quantity of cereals used was large. The quantity of nutrients might easily have been considerably increased, with but little or no increase in the cost of the diet, by the purchase of more cereals and less meat.

#### DIETARY STUDY OF A HOUSEKEEPER'S FAMILY (NO. 168).

This family consisted of the mother, 55 years of age, weighing 130 pounds; her three daughters, one 22, one 20, and the other 14 years old, and weighing 140, 130, and 98 pounds, respectively, and one son, 17 years of age, weighing 120 pounds. The mother and three oldest children

were born in Ireland. All were in excellent health and all were wage-earners. The rent of the rooms occupied was \$15 a month. Of this the mother paid \$9 a month by her work as housekeeper or janitor. One room was let for \$1.50 a week. The total income from the children amounted to \$20 a week, which was all turned into the family treasury. Each one carried a lunch from home and spent 5 cents daily for tea or coffee. The home was kept neat and clean and the table was attractive. The family dressed well and appeared to be in good circumstances. No member of the family was familiar with cooking, and all of the pastry was purchased of a baker; the girls, however, were anxious to learn how to cook.

The study began September 20, 1896, and continued ten days. The number of meals taken was as follows:

Al Al	ears.
Three women (90 meals $\times$ 0.8 meal of man), equivalent to	72
Boy, 17 years old (30 meals $\times$ 0.8 meal of man), equivalent to	24
Girl, 14 years old (30 meals $\times$ 0.7 meal of man), equivalent to	21
Woman visitor (4 meals $\times$ 0.8 meal of man), equivalent to	3
Man visitor	4
Total number of meals taken equivalent to.	124

Table 12.—Weights and cost of food and nutrients in dietary study No. 168.

Equivalent to one man forty-one days.

			V	U	
	Cost, nutrients, and fuel value of food per ma per day.				
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.  Beef: Steak, chuck, 1,50 pounds, 15 cents (27); steak,	Cents.	Grams.	Grams.	Grams.	Calories.
sirloin, 3.50 pounds, 56 cents (33); corned rump, 6 pounds, 60 cents (10); stew meat, 2 pounds, 13 cents (23). Mutton: Chops, 3 pounds, 36 cents (46); chops, 2 pounds, 24 cents (44); leg, 8.50 pounds, \$1.10 (47).	7.7	47	55		704
Pork: Ham, smoked, 9 pounds, \$1.20 (65); bacon, 3.50 pounds, 34 cents (59)	3.7	20 6	63		668 24
Eggs, 1.69 pounds, 28 cents (114) Butter, 4.50 pounds, \$1.05 (118) Milk, 17.07 pounds, 63 cents (124)	2.6 1.5	3 1 6	2 42 8	10	30 394 140
Total animal food		83	170	10	1,960
VEGETABLE FOOD.					
Cereals: Biscuit, 6.94 pounds, 35 cents (139); bread, 10.78 pounds, 48 cents (134); coffee cake, 7.81 pounds, 50 cents (144); corn cake, 2.19 pounds, 10 cents (150); currant loaf, 1.25 pounds, 10 cents (147); doughnuts, 3.50 pounds, 15 cents (157); jelly cake, 0.49 pound, 10 cents (146); rolls, milk, 1 pound, 5 cents (163); rolls, water, 6.69 pounds, 35			1		
cents (166); sweet cake, 0.75 pound, 10 cents (148) Sugars, starches, etc.: Sugar, 9.99 pounds, 48 cents	5, 6	39	33	245	1,471
(169); cocoa, 0.50 pound, 20 cents (171)	1.6	1	1	113	476
cents (215); tomato catsup, 2.50 pounds, 28 cents (217). Fruits: Apples, 11.36 pounds, 20 cents (222); lemons,	1.9	7	1	57	272
1 pound, 5 cents (233); prunes, 0.75 pound, 10 cents (237).	.8	1	1	25	116
Total vegetable food	9.9	48	36	440	2, 335
Total food	26.7	131	206	450	4, 295

The results of this study are interesting. Although the family consisted of women and one 17-year-old boy, the average daily diet furnished more than eighth-tenths as much as the commonly accepted standard requires for a man at moderate labor. Judged by the standard, therefore, they are more than they actually required. If the quantity of protein had been reduced a tenth and the quantity of energy a fifth, the ration would have been still as large as is needed theoretically. The fact that the excess of energy was larger than that of protein indicates that the fats and carbohydrates were in excess. One reason for this is found in the large amount of pork eaten. The quantity of bread and pastry was also large. All such food was purchased of a baker, and its cost was much greater than would have been the case had it been made at home. The amount of cake purchased was quite large, and cost considerably more than the bread eaten. One of the least economical purchases was 2½ pounds of tomato catsup, which cost 28 cents and contained almost no nutriment. If the \$1.20 expended for smoked ham had been used to purchase dried legumes or some of the leaner cuts of beef, the nutritive ratio of the diet would have been improved. During the study \$1.80 was expended for tea and coffee. This sum has not been included in the cost of the food. The income of the family was sufficient to maintain them comfortably, and they apparently made very good use of it.

#### DIETARY STUDY OF A CARETAKER'S FAMILY (NO. 170).

This family is typical of a large number in this region, in which the woman is the breadwinner. It consisted of the mother, 25 years old; her mother, 65 years old and her sister, 17 years old; a child 6 and one 3 years old, weighing 160, 130, 127, 35, and 30 pounds, respectively, and apparently in good health. The mother earned \$15 a month cleaning an office. Her sister paid \$3 a week for room and board, and until a short time previous to the study there had been another boarder who paid \$5 a week for room and board. The rent of four rooms was \$14 a month, but unless they succeeded in obtaining another boarder, the family intended to take cheaper rooms. Food was bought in small quantities for cash and no ice was used. The house was kept neat and clean but the kitchen was so dark that no table was spread. There was no visible waste. Judging from the appearance of the family it would be supposed that their circumstances were better than was actually the case.

The study began October 1, 1896, and continued ten days. The number of meals taken was as follows:

	Meals.
Two women (60 meals $\times$ 0.8 meal of man), equivalent to	. 48
Girl, 17 years old (30 meals $\times$ 0.7 meal of a man), equivalent to	. 21
Child, 6 years old (20 meals × 0.5 meal of man), equivalent to	. 10
Child, 3 years old (20 meals × 0.4 meal of man), equivalent to	8
(D=4=1 1 f 1 - 4 - 1 1 - 4 - 1	OF

Total number of meals taken equivalent to 87 Equivalent to one man twenty-nine days.

Table 13.—Weights and cost of food and nutrients in dietary study No. 17

Kinds, amounts, and cost of food for ten days.	Cost, nutrients, and fuel value of food per man per day.				
	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.  Beef: Corned, 2.50 pounds, 13 cents (2). Veal: Shoulder, 1 pound, 8 cents (58). Lamb: Leg, 7 pounds,	Cents.	Grams.	Grams,	Grams.	Calories.
56 cents (42).  Pork: Head, 1 pound, 5 cents (70); sparerib, 2.36 pounds, 15 cents (81); salt, 2.25 pounds, 17 cents	2,6	25	23		316
(72) Fish: Mackerel, salt, 1 pound, 12 cents (99); oysters,	1.3	9	48		483
0.63 pound, 10 cents (101) Eggs, 0.41 pound, 5 cents (114). Butter, 1.82 pounds, 37 cents (118).	1.3	3	3 1 21		40 13 222
Milk, 2.62 pounds, 5 cents (124) Milk, condensed, 6.51 pounds, 45 cents (125)	1.6	9	8	55	35 336
Total animal food	7.9	49	109	57	1,445
Cereals: Bread, rye, 9.70 pounds, 37 cents (136); bread, wheat, 6.62 pounds, 25 cents (134); macaroni, 1 pound, 10 cents (158); rolls, milk, 1.13 pounds, 5 cents (163); rolls, water, 2.82 pounds, 15 cents (166). Sugar, 4.31 pounds, 22 cents (169). Vegetables: Beans, 1.75 pounds, 9 cents (175); cabbage, 2.50 pounds, 2 cents (179); corn, 0.50 pound, 3 cents (184); onions, 1.25 pounds, 3 cents (195); potatoes, Machael (184); onions, 1.25 pounds, 3 cents (195); potatoes, weet, 1.44 pounds, 17 cents (204); potatoes, sweet,	3.2	31	5	181 67	916 275
1.50 pounds, 3 cents (206); tomatoes 8.50 pounds, 13 cents (215); turnips, 2.50 pounds, 3 cents (218). Fruit: Apples, 5 pounds, 10 cents (221)	1.8	14	2	77 9	392 37
Total vegetable food	6.1	45	7	331	1,620
Total food	14, 0	94	116	391	3,065

The quantity of protein and energy in this study was somewhat below the standard for persons at active exercise. The cost was moderate, evincing careful management. The quantity of animal foods was not large, and for the most part such foods were economically purchased. The leg of lamb and the oysters, however, were expensive in proportion to the nutrients furnished. Had the 56 cents spent for the leg of lamb been used to buy more of the corned beef at the price paid, and the 10 cents spent for oysters used to buy more yeal shoulder, the quantity of nutrients in the diet would have been increased by 18 grams of protein and 185 calories of energy per man per day. If, in addition, the 21 cents spent for cabbage, corn, onions, and tomatoes had been used to purchase more potatoes and sweet potatoes, the ration would have been still further increased by 3 grams of protein and 385 calories of energy per man per day. While these changes would not bring the amount of protein in the ration up to the commonly accepted standard, they indicate how, for the same expenditure, more nutriment could have been obtained than was actually the case. Each family must of course determine the extent to which variety shall give way to economy. Had the diet in actual use been increased by one-sixth it would probably have more nearly met the physiological requirements of the family, and even then, at the prices paid, would have cost but 16.7 cents per man per day.

#### DIETARY STUDY OF A TANNER'S FAMILY (NO. 171).

The family is typical of the sober, honest, and industrious poor, who maintain a continual struggle for the bare necessities of life. It consisted of the father, 45 years old; the mother, 45 years old; three boys, one 19, one 14, and one 10 years old, and two girls, one 12 and the other 8 years of age. Their weights were 168, 134, 135, 75, 65, 57, and 51 pounds, respectively. The father worked in a neighboring tannery and earned \$10 per week. The oldest boy had just obtained night work in a flour mill, but at the time he received no wages, and probably would not receive more than \$3 per week at first. The 14-year-old boy was subject to epilepsy and could neither go to school nor work. He could not be left alone in the house, and thus prevented the mother from going out to work. The other children attended school. The mother had been for years a house servant and was an excellent cook. Bread was baked at home, and it is estimated this was at least \$4 a month cheaper than a corresponding amount of bakers' bread. It was the ambition of the mother to be able some day to buy a whole barrel of flour. There was no visible waste. A table was spread and the family sat down together morning and night. The rent paid for four rooms, two light and two dark, was \$9 a month. The kitchen was supplied with hot and cold water.

The study began October 14, 1896, and continued ten days. The number of meals taken was as follows:

M	ears.
Two men	60
Woman (30 meals $\times$ 0.8 meal of man), equivalent to	24
Boy, 14 years old (30 meals $\times$ 0.8 meal of man), equivalent to Girl 12 and boy 10 years old (60 meals $\times$ 0.6 meal of man), equiv-	24
alent to	36
Girl, 8 years old (30 meals $ imes$ 0.5 meal of man), equivalent to	15
Total number of meals taken equivalent to	159

Table 14.—Weights and cost of food and nutrients in dietary study No. 171.

Equivalent to one man fifty-three days.

	Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates	Fuel value.	
ANIMAL FOOD.  Beef: Corned, 2 pounds, 13 cents (3); steak, chuck, 5.50 pounds, 56 cents (27); steak, round, 5 pounds,	Cents.	Grams,	Grams,	Grams.	Calories.	
3.50 pounds, 36 cents (27) steas, 100nd, 5 pounds, 36 cents (30); tripe, 2 pounds, 10 cents (39). Mutton: Leg, 11.13 pounds, 96 cents (47).  Pork: Chops, 3.50 pounds, 35 cents (61); ham, corned, 2 pounds, 15 cents (66); sansage, 2 pounds, 20 cents	4.0	38	30		435	
(74): shoulder, salt, 5.10 pounds, 35 cents (78)	2.0	15	35		387	
fresh, 5 pounds, 30 cents (89). Eggs, 4.19 pounds, 20 cents (117). Butter, 4.11 pounds, 80 cents (118).	4	9 1	1 30		37 13 279	
Cheese, 1.50 pounds, 20 cents (120)	.4	3 1	4	1	49 15	
Milk, condensed, 2.78 pounds, 17 cents (125)		69	103	13	1,295	

Table 14.—Weights and cost of food and nutrients in dietary study No. 171—Continued.

	Cost, nutrients, and fuel value of food per man						
Finds amounts and out of food for tan days		per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.		
VEGETABLE FOOD.  Cereals; Flour, low grade, 3 pounds, 9 cents (132);	Cents.	Grams,	Grams,	Grams.	Calories,		
flour, prepared, 27.59 pounds, 51 cents (133); bread, 7.91 pounds, 30 cents (134)	1.7	34	4	227 51	1, 108 209		
Vegetables: Greens, 5 pounds, 7 cents (189); onions, 5.56 pounds, 7 cents (195); potatoes, 31.26 pounds, 30 cents (204); tomatoes, 8.50 pounds, 7 cents (215); turnips, 7 pounds, 7 cents (218)	1.1	10	1	64	313		
Total vegetable food	3.3	44	5	342	1,630		
Total food	12.7	113	108	356	2, 925		

The average food consumption per man per day during this study was a trifle below the commonly accepted standard as regards protein. and about one-sixth below it as regards energy. The cost, 12.7 cents per man per day, was small, indicating careful management on the part of the mother. The foods which furnished the least nutriment for the money expended on them were the soup greens, onions, tomatoes, and turnips. While a certain amount of variety in vegetable foods is desirable, a part of the 28 cents thus expended might have been used for the purchase of a larger quantity of potatoes and thus have increased the nutritive value of the ration. The cost of animal foods was 75 per cent of the total cost of the food, which is somewhat larger than usual. In other words, the family expended more than was necessary for meats, which are at best quite expensive. If, for instance, from a third to a half of the 96 cents spent for a leg of mutton had been used to purchase wheat flour, and the rest for a cheaper cut of meat, the quantity of protein and energy per man per day would have been nearer that indicated by the standard for a man at moderate work. If the ration had been increased in amount by about one-tenth it would probably have been sufficient for the needs of the family, and would then have cost but 14 cents per man per day. This family, like those in dietary studies Nos. 159, 161, and 170, manifested a considerable degree of skill in their purchases of food.

#### DIETARY STUDY OF A FOUNDRYMAN'S FAMILY (NO. 172).

This family consisted of the father, 40 years old; his wife, 36 years old; a grandmother, 54 years old; an aunt, 28 years old; four daughters, respectively 14, 11, 4, and 2 years old, and an infant 2 months old. The weights of some members of the family were not reported. The father weighed 167; the mother, 140; the grandmother, 160; the 14-year-old daughter, 79, and the 11-year-old daughter, 70 pounds. The father a foundryman, was industrious. He earned \$10

a week when on full time, but his work was unsteady. At the time of the study the grandmother was bedridden, and the aunt, a rag sorter, was so ill with consumption that she was unable to work. The oldest girl was learning box making. The family rented two rooms, one light and one dark, for which they paid \$5.50 a month. The mother was a good cook and did her own marketing. Owing, however, to their very limited income, food was bought in small quantities, so that she was unable to economize as she might had she been able to purchase larger amounts. The food was of good quality, and there was no waste. Everything about the home was neat and clean and the children were tidy and respectable in appearance.

The study began October 14, 1896, and continued ten days. The number of meals taken was as follows:

N.	rears.
Man	30
Three women (90 meals $\times$ 0.8 meal of man), equivalent to	72
Girl, 14 years old (30 meals $\times$ 0.7 meal of man), equivalent to	21
Girl, 11 years old (30 meals $\times$ 0.6 meal of man), equivalent to	18
Two girls, 4 and 2 years old (60 meals $\times$ 0.4 meal of man), equiva-	
lent to	24
Infant, equivalent to	9
Total number of meals taken equivalent to	174
Equivalent to one man fifty-eight days.	

Table 15.— Weights and cost of food and nutrients in dietary study No. 172.

Kinds, amounts, and cost of food for ten days,	Cost, nutrients, and fuel value of food per ma per day.					
Think, which to provide the capture of the capture	Cost.	Protein.	Fat.	Carbohy-drates.	Fuel value.	
ANIMAL FOOD.  Beef: Neck, 2 pounds, 16 cents (21); liver, 2 pounds, 16 cents (18); shank, hind, 2 pounds, 8 cents (24).	Cents.	Grams.	Grams,	Grams.	Calories.	
Mutton: Leg, 10.50 pounds, 76 cents (48); neck, 2 pounds, 9 cents (49).  Pork: Bacon, 0.50 pound, 6 cents (59); chops, 9.75	2.2	21	18		25	
pounds, 98 cents (61); feet, 5.86 pounds, 35 cents (62); salt, 4 pounds, 39 cents (72)	3.1	18	60		63	
Fish: Blue, 4.50 pounds, 27 cents (84); cod, fresh, 5.61 pounds, 35 cents (89). Butter, 5.75 pounds, \$1.26 (118). Milk, 8.58 pounds, 20 cents (124)		9   1   2	1 38 3	3	$\frac{4}{35}$	
Milk, condensed, 5. 87 pounds, 59 cents (125)		4	4	25	15	
Total animal food	9.8	55	124	28	1,49	
VEGETABLE FOOD.						
Cereals: Flour, 44 pounds, \$1.06 (131); bread, white, 1.1 pounds, 5 cents (134); rolls, 1 pound, 5 cents (166). Sugar, 15, 12 pounds, 70 cents (169). Vegetables: Cabbage, 6.75 pounds, 7 cents (179); potatoes, 29.60 pounds, 31 cents (204); tomatoes,	2. 0 1. 2	40	4	266 118	1, 29 48	
4.50 pounds, 4 cents (215); turnips, 6.37 pounds, 3 cents (219)	.8	7		50	23	
Total vegetable food	4.0	47	4	434	2,0	
Total food	13.8	102	128	462	3,5	

The quantity of energy per man per day in this study was equal to that called for by the commonly accepted standard. The quantity of protein, however, was rather small, although as large as has been found in a considerable number of studies of families of mechanics, farmers, and other working people in comfortable circumstances. The cost, 13.8 cents per man per day, was very reasonable. No suggestion for improvement seems to be called for in this case. The satisfactory results obtained are doubtless due to the fact that the woman was a good cook and could do her own marketing. They are interesting as showing that it was possible under the given conditions to live in New York City on a ration of considerable variety and at the same time of small cost.

The quantities of protein and energy obtained for 10 cents in some of the more important food materials purchased by this family are shown in the following table:

Table 16.—Cost of food materials per pound, and amounts of protein and energy obtained for 10 cents in each, in dietary study No. 172.

		In 1 p	ound.	Amounts	bought fo	r 10 cents.	Total amount
	Price per pound.	Protein.	Fuel value.	Total food.	Protein.	Fuel value.	ex- pended during study.
Beef:	Cents.	Downel	Calarico	Pounds.	Pound.	Calories,	Cents.
Neck			770	1, 25	0.18	970	16
Liver	8.0	. 207	605 -		. 26	720	16
Shank		. 096	405	2,50	. 24	1,005	8
Mutton:	4.			20			
Leg	7.2	. 151	900	1.39	. 21	1.245	76
Neck		. 123	985	2.20	. 27	2, 180	9
Pork:							
Baeon	12.0	.091	2,795	. 83	. 08	2,330	6
Chops	10.0	. 134	1,270	1.00	. 13	1,265	98
Feet	6.0	. 158	1,405	1.67	. 26	2,350	35
Salt	9, 8	.019	3,670	1.02	.02	3,765	39
Fish:							
Bluefish						. 350	27
Fresh cod		. 165	325	1.61	. 18	460	35
Butter	21.9	.010	3,605	. 46		1,615	126
Milk	* 2.3	. 033	325	4.35	. 14	1,385	20
Milk, condensed		. 088	1,520	1,00	. 09	1,515	59
Flour		. 112	1,645	4.17	. 46	6,820	106°
Bread			1,215	2.17	. 20	2,640	5
Rolls, water		. 090	1,300	1.89	. 17	2,440	5
Sugar	4.6		1,860	2.17		9,115	70
Green vegetables: Cabbage,							
tomatoes, turnips				12.50	.15	1,620	14
Potatoes	1.1	. 022	385	9.10	. 21	3,695	31
							_

The food for which they spent the most was butter, which furnished no protein and not so large an amount of energy for a given sum as some of the other food materials. For instance, \$1.26 spent for butter furnished some 21,000 calories of energy, while \$1.06 expended for flour purchased nearly 70,000 calories and in addition over a pound of protein. It may be questioned, however, whether any reduction in the butter would be desirable under the circumstances. The comparison is made simply to show the relative returns for a given sum expended for different foods. The amount of green vegetables used was not large. Such food in reasonable amounts is very useful to give variety to the diet. The meats were all purchased at very low prices, bacon being the only one which cost more than 10 cents a

pound. The mutton neck, at  $4\frac{1}{2}$  cents a pound, furnished a large amount of protein and energy for the money expended, and was by far the most economical of the meats purchased. The fish furnished considerable protein, but not much energy. This was probably a wise purchase, however, since the general tendency is to purchase foods containing relatively too much fuel ingredients as compared with the amounts of protein. The use of fish counterbalanced somewhat the comparatively large amounts of sugar and butter in the diet.

## DIETARY STUDY OF A TRUCKMAN'S FAMILY (NO. 177).

This family consisted of the father, 33 years old; the mother, 29 years old; 3 sons, one 11, one 7, and one 5 years of age, and a male relative 30 years old. The members of the family weighed 130, 149, 63, 50, 40, and 150 pounds, respectively. The father, a truck driver, earned \$2 during the ten days covered by the study. The mother was employed as housekeeper, or janitor, receiving for her services the rent of their rooms and \$4 a month in addition. The cousin was not working on full time, but paid practically all his earnings (\$16 per month) into the family treasury. The two older boys obtained their dinner at school, and the youngest at a kindergarten.

The study began January 26, 1897, and continued ten days. The number of meals taken was as follows:

P	rears.
Two men	60
Woman (30 meals × 0.8 meal of man), equivalent to	24
Boy, 11 years old (22 meals $\times$ 0.6 meal of man), equivalent to	13
Boy, 7 years old (26 meals $\times$ 0.5 meal of man), equivalent to	. 13
Boy, 5 years old (22 meals $\times$ 0.4 meal of man), equivalent to	9
Visitor	. 1
(Patal namelan of models talon agriculant to	100

Table 17.— Weights and cost of food and nutrients in dietary study No. 177.

	Cost, nutrients, and fuel value of food per man per day.						
Kinds, amounts, and cost of food for ten days.		Protein.	Fat.	Carbohy- drates.	Fuel value.		
eef: Flank, skirt steak, 4 pounds, 28 cents (35);	Cents.	Grams.	Grams.	Grams.	Calories.		
neck, 1.20 pounds, 7 cents (21); rib, corned, 5 pounds, 33 cents (9); shank, fore, 6 pounds, 30 cents (23); steak, chuck, 2.50 pounds, 25 cents (27); tripe, pickled, 2 pounds, 10 cents (38); corned, canned, 1 pound, 15 cents (5). Mutton: Chops, 2.50 pounds, 35 cents (45); shoulder, 1 pound, 8 cents							
(51)ork: Ham, boiled, 0.56 pound, 10 cents (64); lard,	4.8	45	46		61		
0.25 pound, 2 cents (69)		1	7		6		
ish: Herring, fresh, 2 pounds, 10 cents (95)		5	1		5		
ggs, 0.63 pound, 10 cents (114)		1	1		]		
utter, 1.68 pounds, 34 cents (118)			16		15		
Iilk, 4.63 pounds, 10 cents (124)	.2	2	2	3 12			
illa, contienden, 2 poundo, 10 cents (120)	.0	4		12			
	7.1	56	75	15	99		

Table 17.—Weights and cost of food and nutrients in dietary study No. 177—Continued.

	Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy-drates.	Fuel value.	
VEGETABLE FOOD.  Cereals: Barley, 0.63 pound, 4 cents (126); flour, 1 pound, 4 cents (131); oatmeal, 2 pounds, 7 cents	Cents.	Grams.	Grams.	Grams.	Calories.	
(128); bread, 25.67 pounds, 96 cents (134); crackers, soda, 1 pound, 5 cents (156); rolls, plain, 4.50 pounds, 20 cents (164). Sugar, 5.25 pounds, 24 cents (169). Vegetables: Cabbage, 4 pounds, 5 cents (179); onions, 2.64 pounds, 8 cents (196); potatoes, 15.30 pounds,	3.4	39	9	223 59	1, 155 245	
20 cents (204); soup greens, 1 pound, 6 cents (189); tomatoes, canned, 4 pounds, 26 cents (216)Fruits: Apples, dried, 1 pound, 10 cents (220)	1.6	6	1	40 8	19	
Total vegetable food	5.9	45	10	322	1,630	
Total food	13.0	101	85	345	2, 62	

It seems hardly probable that this family was properly nourished. The quantities of protein and especially energy per man per day found in the dietary study are considerably below the commonly accepted standards. The prices paid for the animal food were reasonable, and the relative expenditure for animal food—55 per cent of the total—was rather less than the average. The most expensive food material used was canned tomatoes, for which the family expended 5 per cent of the total outlay for food, although the protein obtained was but 0.5 per cent and the energy but 0.4 per cent of the total protein and energy, respectively, in the food. The same outlay would have given a more nutritious diet if a different selection of food had been made in some cases. If the cabbage, onions, soup greens, and tomatoes had been left out of the diet and two-thirds of the money expended for them used to buy dried peas or beans, and the remainder for the purchase of more potatoes and bread, the protein could have been increased to 125 grams and the energy to 3,500 calories per man per day without increasing the cost.

# DIETARY STUDY OF A LONGSHOREMAN'S FAMILY (NO. 178).

This study was made with a family which consisted of the father, 42 years old; the mother, 32 years old; the grandmother, 64 years old; four sons—one 14, one 13, one 9, and one 4 years of age—and an infant 6 months old. The weights of the members of the family, aside from the infant, were 180, 145, 170, 110, 75, 60, and 30 pounds, respectively. The father, a longshoreman, earned \$3 during the ten days covered by the study. The oldest boy received \$3.50 per week in a printing office. The family occupied four small rooms, only one of which was well lighted. The rent was paid by the mother's work as housekeeper or janitor.

The study began February 6, 1897, and continued ten days. The number of meals taken was as follows:

	Meals.
Man	30
Two women (60 meals $\times$ 0.8 meal of man), equivalent to	48
Boy, 14 years old (30 meals $\times$ 0.8 meal of man), equivalent to	. 24
Boy, 13 years old (30 meals $\times$ 0.6 meal of man), equivalent to	18
Boy, 9 years old (30 meals $\times$ 0.5 meal of man), equivalent to	. 15
Boy, 4 years old (30 meals $\times$ 0.4 meal of man), equivalent to	12
Infant, equivalent to	9
Visitor	1

Table 18.—Weights and cost of food and nutrients in dietary study No. 178.

Kinds, amounts, and cost of food for ten days.		Cost, nutrients, and fuel value of food per man				
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy-drates.	Fuel value.	
ANIMAL FOOD.  Beef: Corned, 9.40 pounds, 49 cents (4); shank, hind, 4.31 pounds, 20 cents (24); steak, chuck, 2.75 pounds, 30 cents (27); tripe, pickled, 2 pounds, 10	Cents.	Grams.	Grams.	Grams.	Calories.	
cents (38)	2.1	25	25		335	
1.50 pounds, 12 cents (82). Fish: Cod, salt 15 pound, 8 cents (91); herring, fresh,	1.0	9	23		251 43	
2.76 pounds, 15 cents (95)	.1	1	18		468 168	
Butter, 2.50 pounds, 46 cents (118)	.5	2	3	3	49	
Total animal food	5, 0	43	71	3	850	
VEGETABLE FOOD.  Cereals: Flour, 4 pounds, 11 cents (131): bread, 24.06 pounds, 51 cents (138); buns, 6.88 pounds, 30 cents (140).  Sugar, 6 pounds, 34 cents (169).  Vegetables: Beans, dried, 1.87 pounds, 7 cents (175);	1.8	32	7	171 52	897 213	
cabbage, 5.25 pounds, 7 cents (179); carrots, 0.37 pound, 1 cent (182); onions, 0.66 pound, 3 cents (196); peas, 1 pound, 2 cents (200); potatoes, 22.74 pounds, 30 cents (204); turnips, 4.75 pounds, 6 cents (218).	1.1	11	1	59	295	
Total vegetable food	3.5	43	8	282	1,405	
Total food	8.5	86	79	285	2, 255	

The income of this family during the time of the study was very limited, indeed, and the amount expended for food, 8.5 cents per man per day, was unusually small. The food materials obtained for this sum supplied 86 grams of protein and 2,255 calories of energy, which gives evidence of considerable careful management on the part of the mother. The quantity of nutrients, however, was insufficient for the proper nourishment of the family. It is interesting to note that in this case 2 pounds of dried beans were used during the time of the study. The prices paid for meats were very reasonable and the cost of vegetables, other than potatoes, was small.

The following table shows the quantity of protein and energy in 10 cents' worth of the different food materials purchased by this family:

Table 19.—Cost of food materials per pound, and amounts of protein and energy obtained for 10 cents in each, in dietary study No. 178.

		In 1 pou		Amounts	bought for	r 10 cents.	Total
Kind of food material.	Price per pound.	Protein.	Fuel value.	Total food.	Protein.	Fuel value.	amount ex- pended during study.
Beef:	Cents.	Pound.	Calories.	Pounds.	Pound.	Calories.	Cents.
Corned	5.2	0.183	1,385	1,92	0.35	2,650	49
Hind shank	4.6	. 096	405	2.17	.21	870	20
Chuck steak		.166	735	.92	.15	675	30
Pickled tripe		. 117	270	2.00	. 23	545	10
Pork:							
Loin	. 7.7	. 166	1,580	1.30	. 21	2,045	42
Trimmings	8.0	. 050	2,835	1.25	. 06	3,535	12
Fish:							
Salt cod	8.0	.190	3,670	1.25	. 24	460	8
Fresh herring		. 195	660	1.85	. 36	1,215	15
Butter		. 010	3,605	. 54		1,960	. 46
Milk		. 033	325	3.13	.10	1,020	24
Flour	2.8	.112	1,645	3.57	. 41	5, 975	11
Bread	2.1	. 109	1, 215	4.76	. 51	5, 920	51
Buns		. 081	1,450	2.27	.19	3,325	30
Sugar	5.7		1,860	1.75		3, 280	34
Beans, dried	. 3.7	, 225	1,605	2.70	. 60	4, 285	7
Green vegetables: Cabbage,							
carrots, onions, peas, tur-	1 0			6, 25	01	1 055	19
nips		. 022	385	7, 69	.21	1,855 2,945	30
Lutatues	1.0	. 022	220	7.09	.11	2,940	50

It will be seen that flour, stale bread, and beans were the most economical sources of both protein and energy. As sources of protein, the pork trimmings and the milk, and as sources of energy some of the meats were the least economical. The most economical meat was the corned beef, which the family obtained for about 5 cents a pound; the most expensive was the chuck steak. It is difficult to make suggestions for improvement in such a case unless more money was available for the purchase of food. If the family had been willing to eat more beans and bread, less vegetables, less meats, and rather less sugar, more nutriment could have been obtained for the same money. If they had had means to increase the ration one-third in amount with the same relative distribution of purchases as was actually found, the protein and energy would have been nearly sufficient for their needs, and the cost would have been less than 12 cents per man per day.

### DIETARY STUDY OF A CARPENTER'S FAMILY (NO. 180).

This study was made with the same family as in studies No. 166 above, and No. 31 previously published." The father had work at his trade of carpentering only a few hours a day, and earned not more than \$5 a week. The older son (aged 20 years) gave his mother \$7 a week. The 14-year-old daughter also gave her wages of \$3 a week to her mother. The family were in good health.

The study began February 17, 1897, and continued ten days. The number of meals taken was as follows:

	DICHIS.
Two men	_ 60
Woman (30 meals × 0.8 meal of man), equivalent to	24
Girl, 14 years old (30 meals $\times$ 0.7 meal of man), equivalent to	. 21
Boy, 11 years old (30 meals $\times$ 0.6 meal of man), equivalent to	. 18
Total number of meals taken equivalent to.	123
•	
Equivalent to one man forty-one days.	

Table 20.— Weights and cost of food and nutrients in dictary study No. 180.

Kinds, amounts, and cost of food for ten days.		utrients, a	nd fuel va per day.	lue of food per man		
		Protein.	Fat.	Carbohy- drates.	Fuel value.	
ANIMAL FOOD.  Beef: Shoulder, 2.50 pounds, 25 cents (26); steak, round, 5.06 pounds, 65 cents (29); fore shank, 1	Cents.	Grams.	Grams.	Grams.	Calories.	
pound, 7 cents (23); kidney, 2.29 pounds, 20 cents (17); cottolene, 2.13 pounds, 22 cents (12).  Pork: Loin, fresh, 2.81 pounds, 33 cents (81); ham,	3.4	20	35		406	
smoked, 2.76 pounds, 50 cents (65); ham, boiled, 0.80 pound, 10 cents (64). Chickens, 3.50 pounds, 40 cents (83). Eggs, 6.12 pounds, 71 cents (117).			23 6 6		263 84 89	
Butter, 3.31 pounds, 77 cents (118) Cheese, 0.19 pound, 5 cents (120). Milk, 37.75 pounds, 90 cents (124)		1		21	258 13 302	
Total animal food	12.6	62	119	21	1, 445	
VEGETABLE FOOD.						
Cereals: Flour, 4.82 pounds, 15 cents (131); rice, 1.50 pounds, 8 cents (130); prepared flour, 2.13 pounds, 8 cents (133); bread, 17.68 pounds, 56 cents (134); bread, brown, 0.94 pound, 2 cents (135); rolls, plain, 11.62 pounds, 60 cents (164); doughnuts, 3.37 pounds, 20 cents (157). Sugar, 3.93 pounds, 15 cents (169) Vegetables: Beans, 1.75 pounds, 7 cents (175); cab-	4.1	44	17	276 44	1,470 180	
bage, 3.50 pounds, 5 cents (179); onions, 1.19 pounds, 3 cents (196); peas, green, 2 pounds, 5 cents (199); potatoes, 33.31 pounds, 39 cents (201); soup greens, 0.68 pound, 5 cents (189); tomatoes, 1 pound, 8 cents (215); sauerkraut, 3.63 pounds, 10 cents (212)	2.0	15	1	87	428	
Fruits: Apples, 3.92 pounds, 7 cents (222); raspberry jelly, 1 pound, 14 cents (232)	.5		1	14	67	
Total vegetable food	7.0	59	19	421	2, 145	
Total food	19.6	121	138	412	3,590	

The quantities of protein and energy per man per day found in this dietary study were in close agreement with those called for by the commonly accepted standard for men at moderate muscular work. The cost, while reasonable, was considerably larger than in some of the previous studies, and in view of the small and uncertain income of the father might advantageously have been reduced by the substitution of cereal foods and dried legumes for the larger portion of the vegetables other than potatoes, and for part of the meats. It must be remembered that while meats form an appetizing part of the diet, and one which the average American workman thinks he can not do with out, they are probably not absolutely indispensable.

### DIETARY STUDY OF A SAIL RIGGER'S FAMILY (NO. 183).

This study was made in a family comprising the father, 50 years old, Irish born; the mother, 45 years old, also Irish born; and three sons, one 15, one 12, and one 6 years of age, weighing 200, 180, 78, 60, and 40 pounds, respectively. They were all in good health. The father, a sail rigger, earned \$3 per day when working full time, but seldom was fully employed. During the time covered by the study he worked but two days on account of wet weather. Three rooms, all light, cost the family \$10 a month.

The study began March 13, 1897, and continued ten days. The number of meals taken was as follows:

716	eus.
Man	30
Woman (30 meals × 0.8 meal of man), equivalent to	24
Boy, 15 years old (30 meals $\times$ 0.8 meal of man), equivalent to	24
Boy, 12 years old (30 meals × 0.6 meal of man), equivalent to	18
Boy, 6 years old (24 meals $\times$ 0.5 meal of man), equivalent to	12
_	
Total number of meals taken equivalent to	108
Equivalent to one man thirty-six days.	

Table 21.—Weights and cost of food and nutrients in dietary study No. 183.

	Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.	
ANIMAL FOOD.	Cents.	Grams.	Grams.	Grams.	Calories.	
Beef: Flank, corned, 4.68 pounds, 35 cents (44a); heart, 3.88 pounds, 12 cents (16); neck, 2 pounds, 15 cents (21); steak, chuck, 3.25 pounds, 41 cents (28) Pork: Bacon, 1.24 pounds, 10 cents (59); chops, 2.32 pounds, 35 cents (60); feet, pickled, 8.37 pounds,	2.8	27	42		501	
42 cents (63)	2.4	23	34		411	
Fish: Cod, fresh, 2.81 pounds, 21 cents (90)	.6	6 2	1		25 18	
Eggs, 1.14 pounds, 13 cents (117). Butter, 0.62 pound, 13 cents (118). Milk, 11.12 pounds, 33 cents (124).	.4	5	$\frac{1}{7}$	7	65 105	
Total animal food.	7.5	63	90	7	1,125	
VEGETABLE FOOD.						
Cereals: Barley, pearled, 0.31 pound, 2 cents (126); bread, 40.36 pounds, \$1.32 (134). Sugar, 5.75 pounds, 29 cents (169). Vegetables: Beans, 2 pounds, 8 cents (175); cabbage, 8.99 pounds, 20 cents (180); onions, 1.25 pounds, 6 cents (196); potatoes, 28.75 pounds, 45 cents (204);	3.7	47	7	273 73	1,376 299	
soup greens, 1.25 pounds, 5 cents (189); tomatoes, 4 pounds, 16 cents (216); turnips, 1.75 pounds, 2 cents (218).	2,9	17	1	93	460	
Total vegetable food	7.4	64	8	439	2,135	
Total food	14.9	127	98	446	3, 260	

The quantity of protein consumed per man per day by this family was slightly larger, while the energy was somewhat smaller than called for by the commonly accepted standard for a man at moderate work.

Inasmuch as protein is the most expensive nutrient and is furnished by animal foods in relatively larger proportions than in vegetable foods, it would appear that this family might have had a rather better balanced ration by substituting cereals for a portion of the animal food. Such a change would certainly not have increased and perhaps might have diminished the cost of the diet. As it was, however, the cost was very reasonable, amounting to but 15 cents per man per day.

## DIETARY STUDY OF A DAY LABORER'S FAMILY (NO. 185).

This family comprised the father, 45 years old; the mother, 45 years old, and four children; two boys, one 21 and the other 16 years old, and two girls, one aged 13 and the other 7 years. Their weights were 169, 235, 169, 130, 89, and 63 pounds, respectively. The health of the family was good. The father, a day laborer, had been idle for some time. The mother earned from \$1 to \$1.50 a week washing. The young man was a helper on a truck wagon receiving \$3 a week wages. The rent of two back rooms, one dark, which they occupied was \$5.50 a month. Food was bought by the meal and there was no visible waste. They used stale bread which could be purchased at half the price of fresh bread. Any food left from one meal remained on the table till the next.

The study began March 2, 1897, and continued ten days. The number of meals taken was as follows:

	Mea	als.
Two men		60
Woman (30 meals $\times$ 0.8 meal of man), equivalent to		24
Boy, 16 years old (30 meals $\times$ 0.8 meal of man), equivalent to		24
Girl, 13 years old (30 meals $\times$ 0.6 meal of man), equivalent to		18
Boy, 7 years old (30 meals $\times$ 0.5 meal of man), equivalent to		15
Total number of meals taken equivalent to	1	41
Equivalent to one man forty-seven days.		

Table 22.—Weights and cost of food and nutrients in dietary study No. 185.

Kinds, amounts, and cost of food for ten days,	Cost, nutrients, and fuel value of food per man per day.						
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.		
ANIMAL FOOD.  Beef: Shoulder clod, 2 pounds, 20 cents (26); skirt-	Cents.	Grams.	Grams.	Grams.	Calories.		
ing steak, 3.50 pounds, 36 cents (36)  Pork: Shoulder, corned, 5.56 pounds, 42 cents (78):	1.2	9	8		112		
sausage meat, 2 pounds, 18 cents (75) Eggs, 1.12 pounds, 25 cents (114)	1.3	12 1	24 1		273 14		
Butter, 2.75 pounds, 54 cents (118)	1.1	3	23 3		. 214		
Milk, 18.70 pounds, 46 cents (124)	1.0	6	7	9	127		
Total animal food	5.4	31	66	9	780		

Table 22.—Weights and cost of food and nutrients in dietary study No. 185—Continued.

		Cost, nutrients, and fuel value of food per man per day.						
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.			
VEGETABLE FOOD.  Cereals: Oatmeal, 5 pounds, 20 cents (128); bread, 39.37 pounds, 78 cents (138); buns, 1.14 pounds, 5	Cents.	Grams.	Grams.	Grams.	Calories.			
cents (140); crackers, soda, 0.50 pound, 4 cents (156); cake, 0.50 pound, 6 cents (142) Sugars, starches, etc.: Molasses, 6.20 pounds, 22 cents	2.4	51	10	249	1,323			
(170); sugar, 4.56 pounds, 24 cents (169); cornstarch, 0.37 pound, 3 cents (172) Vegetables: Cabbage, 4 pounds, 4 cents (180); potatoes, 9.19 pounds, 21 cents (204); turnips, 3.44	1.0	1		89	369			
pounds, 7 cents (218)	.7	3		21	98			
Total vegetable food	4.1	55	10	359	1,790			
Total food	9, 5	86	76	- 368	2,570			

The family here studied, like that in No. 178, had a very small income. This fact was apparent in the quantity of nutrients and energy in the food eaten. A considerable degree of good management was displayed, however, in the selection of food, which cost but 9.5 cents per man per day. The chief suggestion for the improvement of this dietary would be an increase in the amounts of protein and energy by one-fourth or two-fifths. If this were done by increasing proportionally the quantities of food materials actually used it would make the cost only about 11.5 cents per man per day. If, however, the family felt that they could not increase the living expenses, an increase of nutriment might still have been obtained by using less animal food and a corresponding larger amount of cereal foods. This might, of course, have detracted to some extent from the palatability of the diet according to the opinion of the average working man, but would have furnished the protein and energy required for proper nourishment of the body. At the same time the diet would have been wholesome.

# DIETARY STUDY OF A FRUIT VENDER'S FAMILY (NO. 186).

This study was carried on in an Italian family comprising the husband, 55 years; his wife, 36 years; a nephew of 16 years, a niece of 8 years, a brother-in-law, 45 years, and his wife, 40 years of age. The weights of the members of the family were 200, 130, 120, 45, 120, and 130 pounds, respectively. All were natives of Italy, and the three men were employed at fruit vending. The head of the family earned \$10 or \$12 a week, and his wife, who sewed for a clothing house, earned a few dollars a week. The brother-in-law and nephew worked for their board. During the study the men worked eight days. They had a cup of coffee when they went out to work at 4 o'clock in the morning, and later in the day purchased a cup of coffee and some rolls; the other meals were taken at home. It has been assumed that,

as the food eaten away from home was small in amount, each one of the men had the equivalent of twenty-six full meals at home during the study. Ten dollars a month rent was paid for the three rooms which they occupied. Food was bought by the day.

The study began March 17, 1897, and continued ten days. The number of meals taken was as follows:

71	eals.
Two men	52
Two women (60 meals $\times$ 0.8 meal of man), equivalent to	48
Boy, 16 years old (26 meals × 0.8 meal of man), equivalent to	21
Girl, 8 years old (30 meals $\times$ 0.5 meal of man), equivalent to	15
Total number of meals taken equivalent to	136

Table 23.—Weights and cost of food and nutrients in dictary study No. 186.

Equivalent to one man forty-five days.

		Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy-drates.	Fuel value.		
ANIMAL FOOD.  Beef: Chopped, 1.50 pounds, 15 cents (29); shank, fore, 6 pounds, 44 cents (23); steak, sirloin, 4 pounds, 52 cents (33); steak, round, 2 pounds, 24 cents (29); suet, 1 pound, 5 cents (37). Veal: Chops.	Cents.	Grams.	Grams.	Grams.	Calorics.		
1.61 pounds, 20 cents (54); cutlets, 1 pound, 14 cents (55)		27   5   5   12			353 393 142		
Fish: Cod, salf, 3 pounds, 18 cents (91); shad, fresh, 2.81 pounds, 28 cents (108). Eggs, 12.12 pounds, \$1.29 (114). Butter, 2.74 pounds, 55 cents (118). Cheese, 2 pounds, 34 cents (129). Milk, 40.98 pounds, 96 cents (124)	2.9 1.2 .7	11 16 5 14	3 13 23 7 17	21	73 187 214 86 302		
Total animal food	15.2	90	139	21	1,750		
VEGETABLE FOOD.  Cereals: Corn meal, 2 pounds, 8 cents (151); rice, 4 pounds, 25 cents (130); macaroni, 6.50 pounds, 40 cents (158); vernicelli, 1 pound, 9 cents (168); bread, 25.08 pounds, \$1.16 (134); cake, 1.25 pounds, 15 cents (142); pie, apple, 1 pound, 10 cents (160). Sugars, starches, oils, etc.: Sugar, 6 pounds, 29 cents, (169); cocoa, 0.50 pound, 11 cents (171); olive oil, 1.36 pounds, 34 cents (173)  Vegetables: Beans, dry, 1 pound, 5 cents (175); cabbage, 4 pounds, 6 cents (180); greens, dandelion, 2 pounds, 17 cents (188); greens, 5.43 pounds, 30 cents (189); onions, 1 pound, 5 cents (195); parsley, 0.12 pound, 1 cent (197); pickles, mixed, 1 pound, 10 cents (203); potatoes, 10.49 pounds, 14 cents (204); spinach, 4.12 pounds, 33 cents (214); tomatoes, 8 pounds, 30 cents (215).	5.0	39 1	7 15	249 62 39	1, 246 397		
Fruits: Jelly, currant, 0.75 pound, 8 cents (232); nuts, 0.50 pound, 5 cents (242)	3	1	2	6	47		
Total vegetable food	9.3	51	25	356	1,900		
Total food	24.5	141	164	377	3,650		

Judged by the usual dietary standards, this family consumed food in excess of their needs. The protein might have been reduced by one-seventh and still have been sufficient according to the commonly accepted dietary standards. The energy, however, could have been reduced but very little. The cost of the food, 24.5 cents per man per day, was much greater than in some of the dietary studies previously described. This was due in part to the use of more expensive meats and in part to the purchase of green vegetables. The cost of the diet could have been largely reduced by the selection of cheaper meats and fish, the use of fewer eggs, and especially by decreasing the amount of green vegetables and using cereals and dry legumes in their place. The relative values of some of the principal items in this study are illustrated in the following table, showing the amounts of protein and energy in 10 cents' worth of each at prices actually paid per pound:

Table 24.—Cost of food materials per pound, and amounts of protein and energy obtained for 10 cents in each, in dietary study No. 186.

	Drice non	In 1 pe	ound.	Amounts	bought fo	r 10 cents.	Total amount
Kind of food material.	Price per pound.	Protein.	Fuel value.	Total food.	Protein.	Fuel value.	ex- pended during study.
Beef:	Cents.	Pound.	Calories.	Pounds.	Pound.	Calories.	Cents.
Chopped	10.0	0.190	895	1.00	0.19	895	15
Fore shank	7.3	.128	545	1.37	. 18	745	44
Sirloin	13.0	. 165	985	.77	. 13	760	52
Round	12.0	. 190	895	. 83	. 16	745	24
Suet	5.0	.047	3,540	2,00	. 09	7,080	5
Veal:	12.5	.199	825	. 80	. 16	660	20
Chops	14.0	.199	690	71	. 16	495	14
Pork:	14.0	. 201	050	. / 1	. 14	450	1.1
Chops	9.6	. 166	1,580	1.04	. 17	1,645	26
Ham	20.0	. 142	1,675	, 50	. 07	835	10
Lard	6.0		4, 220	1,67		7,035	18
Chicken	16.0	. 193	1,045	. 63	. 12	655	98
od, salt	6.0	. 190	315	1, 67	. 32	525	18
had	10.0	. 188	750	1.00	19	750	28
Eggs	10.6	. 134	720	. 94	. 13	680	129
Butter	20.0	. 010	3,605	. 50		1,800	55
Cheese	17.0	. 259	1,950	. 59	. 15	1,145	34
dilk	2, 3	. 033	325	4.35	14	1,415	96
Corn meal	4.0	. 092	1,655	2.50	. 23	4,140	8
Rice	6.2	. 080	1,630	1, 61	. 13	2,630	25
Jacaroni	6.2	. 134	1,665	1.61	. 22	2,685	40
Vermicelli	9.0	. 109	1,625	1.11	. 12	1,805	9
Bread	4.6	. 092	1,215	2.17	. 20	2,640	116
Cake	12.0	. 063	1,675	. 83	. 05	1,395	15
Pie, apple	10.0	. 031	1,270	1.00	. 03	1,270	10
ugar	4.8	(110	1,860	2.08	10	3,875	29 11
Cocoa	22.0	. 216	2,320 4,220	. 45	. 10	1,055	34
Olive oil	25.0	. 225		2.00	. 45	$\frac{1,690}{3,210}$	59
Beans, dried	5. 0 1. 5	. 014	1,605 $125$	6, 67	. 09	835	6
Cabbage	8.5	.024	285	1.18	. 03	335	17
Greens, dandelion	5, 5	. 042	283	1. 18	.08	400	30
onions	5.0	. 014	205	2.00	. 03	410	5
Pickles	10.0	.011	110	1.00	.01	110	10
Potatoes	1.3	.022	385	7.69	.17	2,960	14
Spinach	8.0	. 021	110	1.25	. 03	140	33
Comatoes	3, 8	. 009	105	2.63	. 02	275	30
elly	10.7	.020	1,560	. 93	. 02	1,460	8
Nuts	10.0	.079	1,600	1.00	.08	1,600	5

## DIETARY STUDY OF A WATCHMAN'S FAMILY (NO. 187).

This study was made in the same family as that in dietary study No. 34, reported in a previous publication." The family comprised the father, 50 years old; the mother, 40 years old; an aunt, 26 years old, and seven children—three boys, one 20, one 15, and one 13 years, and four girls, one 16, one 12, one 7, and one 3 years of age.

The weights of the different members were 150, 120, 95, 117, 83, 65, 160, 60, 50, and 30 pounds, respectively. The father, a night watchman, was unemployed at the time of the study; the mother went out cleaning and earned \$3 a week. The oldest son was a printer, and always gave his mother \$10 or \$12 a week. The older daughter was learning to make kid gloves and earned \$3 a week. The aunt was employed at book folding, and paid \$3 a week for board. The family paid \$13 a month rent for four rooms. Food was purchased by the day for cash. The family were thrifty in their habits and neat in appearance.

The study began March 24, 1897, and continued ten days. The number of meals taken was as follows:

	Meals.
Two men	. 60
Two women (56 meals $\times$ 0.8 meal of man), equivalent to	. 45
Girl, 16 years old (30 meals $\times$ 0.7 meal of man), equivalent to	. 21
Boy, 15 years old (30 meals $\times$ 0.8 meal of man), equivalent to	. 24
Boy and girl, 13 and 12 years old (60 meals $\times$ 0.6 meal of man)	,
equivalent to	. 36
Girl, 7 years old (30 meals $\times$ 0.5 meal of man), equivalent to	. 15
Girl, 3 years old (30 meals $\times$ 0.4 meal of man), equivalent to	. 12
Total number of meals taken equivalent to	. 213
Equivalent to one man seventy-one days.	

Table 25.—Weights and cost of food and nutrients in dietary study No. 187.

Vinds amounts and and of find for the day	Cost, nutrients, and fuel value of food per man per day.						
Kinds, amounts, and cost of food for ten days.		Protein.	Fat.	Carbohy- drates.	Fuel value.		
ANIMAL FOOD.  Beef: Steak, sirloin, 4.68 pounds, 60 cents (33); corned, 5 pounds, 30 cents (3). Lamb, 3.5 pounds, 32 cents (43a).  Pork: Ham, 5 pounds, 69 cents (66).  Chicken, 3.37 pounds, 39 cents (83).	Cents.	Grams. 14 4 3	Grams. 19 11 3	Grams.	Calories.  234 118 40		
Fish: Salmon canned, I pound, 18 cents (104); smelts, 5 pounds, 45 cents (109); oysters, 1.38 pounds, 25 cents (101).  Eggs, 7.22 pounds, 63 cents (114).  Butter, 4.62 pounds, \$1.07 (118).  Cheese, 0.5 pound, 8 cents (120).  Milk, 12.07 pounds, 29 cents (124)	1.5 .1 .4	7 6 1 3 3	1 5 25 1 3 3	19	38 71 232 13 56 118		
Total animal food	7.9	41	71	23	920		
VEGETABLE FOOD.  Cereals: Barley, pearled, 0.44 pound, 2 cents (126); bread, 6.25 pounds, 20 cents (134); bread, rye, 39.2 pounds, \$1.42 (136); crackers, soda, 1 pound, 7 cents (156).  Sugar, 13.5 pounds, 62 cents (169).  Vegetables: Beans, 2 pounds, 8 cents (175); cabbage, 2.94 pounds, 8 cents (179); cabbage, 2.74 pounds, 7 cents (181); carrots, 0.62 pound, 1 cent (182); greens, 0.5 pound, 3 cents (189); onions, 0.87 pound, 3 cents (196); peas, dried, 1 pound, 4 cents (200); potatoes,	2.4	27	2	162 86	794 353		
24.95 pounds, 29 cents (204); tomatoes, 2 pounds, 8 cents (215); tomato catsup, 2 pounds, 5 cents (217) Fruits: Prunes, 6.24 pounds, 58 cents (237)	1.2	10 1	1	46 29	240 123		
Total vegetable food	5.2	38	3	323	1,510		
Total food	13.1	79	74	346	2, 43		

The quantities of protein and energy in this study were below those which it is believed suffice for the proper nourishment of a man at moderate muscular work and should have been increased by about 40 per cent. The cost of the increased ration, provided it consisted of the same kinds of food materials and in the same proportions, would have been about 18.5 cents per man per day. Such a sum is quite reasonable. However, the amounts of nutrients might have been increased, without much increase in cost, by the purchase of cheaper meats and fewer green vegetables and by the use of more cereals and legumes.

# DIETARY STUDY OF A BOOKBINDER'S FAMILY (NO. 188).

This study was made in a family which consisted of the father, 36 years old, the mother, 39 years old, and eight children—six daughters, aged, respectively, 18, 16, 14, 10, 6, and 4 years, and two sons, one 12 and the other 2 years of age. The weights of the family were 165, 125, 110, 100, 86, 65, 50, 40, 75, and 35 pounds, respectively. The father, a bookbinder, who had worked in the same place for twenty years, earned \$12 a week. The two older girls were wage-earners, one working at bookbinding, the other at kid-glove making. Each paid their mother \$2.50 a week. This family had occupied the same four rooms, for which they paid \$13 a month rent, for thirteen years. Food was bought in small quantities for cash. There was no avoidable waste.

The study began March 24, 1897, and continued ten days. The number of meals taken was as follows:

	Meals.
Man	. 30
Two women (60 meals × 0.8 meal of man), equivalent to	. 48
Two girls, 16 and 14 years old (60 meals $\times$ 0.7 meal of man), equiv	7-
alent to	. 42
Two children, 12 and 10 years old (60 meals × 0.6 meal of man)	,
equivalent to	. 36
Child, 6 years old (30 meals $\times$ 0.5 meal of man), equivalent to	. 15
Two children, 2 and 4 years old (60 meals × 0.4 meal of man)	,
equivalent to	. 24
Total number of meals taken equivalent to	. 195
Equivalent to one man sixty-five days.	

Table 26.—Weights and cost of food and nutrients in dietary study No. 188.

	Cost, n	utrients, ar	nd fuel va per day.	per man	
Kinds, amounts, and cost of food for ten days.		Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.	Claudo	Cagna	Cuann	Cumus	Calanias
Beef: Corned, 11.50 pounds, \$1.15 (3); steak, round,	Cents.	trrams.	(Trums.	Grams.	Catories.
4 pounds, 48 cents (29); steak, sirloin, 4 pounds, 53 cents (33). Fish: Cod, fresh, 3 pounds, 24 cents, (89); cod, salt,	3, 3	22	. 27		341
3 pounds, 24 cents (91); oysters, 3.30 pounds, 25 cents (101); salmon, canned, 1 pound, 18 cents (104).	. 6	9 4	1 3		50 44
Butter, 5.5 pounds, \$1.65 (118)	. 2	2 12		18	307 26 262
Total animal food	10.5	49	81	19	1,030
VEGETABLE FOOD,				-	
Dereals: Rice, 1 pound, 6 cents (130); bread, rye, 1.06 pounds, 5 cents (136); bread, wheat, 39.32 pounds, \$1.79 (131); buns, 0.88 pound, 5 cents (140); cackes, sweet, 1 pound, 16 cents (142); crackers, 0.50 pound, 9 cents, (152); muffins, 0.68 pound, 5 cents (159); rolls, plain, 3.76 pounds, 20 cents (164)	3.8	31	6	183   60	933 246
Vegetables: Corn, canned, 3 pounds, 20 cents (185); potatoes, 25.12 pounds, 28 cents (204); tomatoes, canned, 8 pounds, 28 cents (216).	1.2	5 1	1		189
Fruits: Prunes, 1.57 pounds, 16 cents (237)	.2			8	32
Total vegetable food	5.8	36	7	290	1,400
Total food	16.3	85	88	309	2, 430

As in the previous study, the quantities of nutrients and energy consumed per man per day should have been about 40 per cent larger in order to equal the amounts usually considered as desirable for men at moderate work. The father, however, worked indoors and at not especially active labor, so that it may be that the family needed rather less than is called for by the standard suggested. At the time of the study the family expended \$7.40 per week for food, which was about half their income. If the diet selected had been increased by 40 per cent the cost would have been 22.8 cents per man per day. The relatively high cost as compared with that observed in some of the previous studies is accounted for by the use of more expensive meats, by the higher price paid for some of the cheaper cuts of meat, and probably also for bakers' goods, and by the expenditures for canned corn and canned tomatoes.

#### DIETARY STUDY OF A BUTCHER'S FAMILY (NO. 192).

The family in this study consisted of the husband (a butcher's assistant) and his wife, both natives of Ireland. The husband was sober and industrious and earned \$11 per week. He also received a late breakfast at his employer's expense. His work began very early in the morning. The family paid \$7.50 a month rent for two rooms. Food was bought by the day for cash.

The study began April 3, 1897, and continued ten days. The number of meals taken was as follows:

MI	ears.
Man	22
Woman (30 meals × 0.8 meal of m a), equivalent to	24
Woman visitor (5 meals $\times$ 0.8 meal of man), equivalent to	4
-	
Total number of meals taken equivalent to	50
Equivalent to one man seventeen days.	

Table 27.—Weights and cost of food and nutrients in dietary study No. 192.

Kinds, amounts, and cost of food for ten days.	Cost, nutrients, and fuel value of food per man per day.					
	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.	
ANIMAL FOOD.  Beef: Steak, sirloin, 2 pounds, 24 cents (33). Lamb	Cents.	Grams.	Grams.	Grams.	Calories.	
chops, 5.5 pounds, 46 cents (46). Mutton, shoulder, 3 pounds, 15 cents (51).  Pork: Ham, 5 pounds, 60 cents (66).  Fish: Cod, 2 pounds, 16 cents (89).	3.5	43   19   6   17	56 45		697 496 24 200	
Eggs, 5.43 pounds, 42 cents (117). Butter, 0.75 pound, 21 cents (118). Milk, 1.25 pounds, 4 cents (124). Milk, condensed, 1 pound, 7 cents (125)	1.2	1 3	17 1 1 2	2 14	158 22 88	
Total animal food	13.8	89	135	16	1,685	
VEGETABLE FOOD.				1		
Cercals: Bread, 8.25 pounds, 29 cents (134) Sugars, starches, and oils: Sugar, 3.75 pounds, 18	1.7	20	3	117	590	
cents (169); cocoa, 0.5 pound, 16 cents (171)	2.0	3	4	105	480	
11.74 pounds, 15 cents (204); tomatoes, 2 pounds, 7 cents (215); turnips, 2.75 pounds, 3 cents (218)	1.9	11	1	74	360	
Total vegetable food	5.6	34	8	296	1,430	
Total food	19.4	123	143	312	3,115	

In this study the cost of food per man per day is not excessive in proportion to the income, while the nutrients and energy obtained are not far from the amounts called for by the usual dietary standard.

# DIETARY STUDY OF A SAIL RIGGER'S FAMILY (NO. 193).

The family here studied consisted of the father, born in Germany, aged 42 years, weighing 160 pounds; the mother, born in Sweden, aged 52 years, weighing 202 pounds; and a daughter, 6 years old, weighing 52 pounds. The father, a sail rigger, earned \$21 per week. On working days he bought his breakfasts at a restaurant. The family occupied four rooms, for which they paid \$14 a month rent, but sublet one room for \$6 a month. Food was bought for cash in small quantities at the small markets.

The study began April 3, 1897, and continued ten days. The number of meals taken was as follows:

Mo	eals.
Man	23
Woman (30 meals × 0.8 meal of man), equivalent to	24
Girl, 6 years old (30 meals $\times$ 0.5 meal of man), equivalent to	
Total number of meals taken equivalent to	62
Equivalent to one man twenty-one days.	

Table 28.—Weights and cost of food and nutrients in dietary study No. 193.

Winds amounts and cost of food for ton days	Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.	
ANIMAL FOOD.  Beef: Round steak, 4.5 pounds, 54 cents (29); blood,	Cents.	Grams.	Grams.	Grams.	Calories.	
2 pounds, 10 cents (39a). Veal: Cutlets, 1.5 pounds, 12 cents (55); leg, 5.5 pounds, 55 cents (56)	6. 2 1. 4	47 9	24 8		416 111	
pounds, 24 cents (103); perch, yellow, 2 pounds, 15 cents (102); sardines, canned, 1 pound, 25 cents (107) 2ggs, 9.37 pounds, 75 cents (114)	3.6 3.6 2.3	20 27 1	3 21 39		110 306 367	
Butter, 2.11 pounds, 49 cents (118) Milk, 7.44 pounds, 19 cents (124) Milk, condensed, 7 pounds, 49 cents (125)	2.3	5 13	7 12	8 82	119 501	
Total animal food	20.3	122	114	90	1,930	
VEGETABLE FOOD.						
Cereals: Flour, 3 pounds, 12 cents (131); cake, coffee, 1.13 pounds, 8 cents (143); bread, rye, 3.88 pounds, 18 cents (136); bread, wheat, 2 pounds, 8 cents (134); rolls, plain, 1 pound, 5 cents (164).  Sugar, 4.87 pounds, 20 cents (169).  Vegetables: Leeks, 0.39 pound, 1 cent (192); onions, 1 pound, 2 cents (195); parsnips, 4.25 pounds, 2 cents (198); potatoes, 17.92 pounds, 20 cents (204); sweet	2.4	23	5	144 105	731 431	
potatoes, 0.36 pound, 1 cent (207); tomatoes, 6 pounds, 21 cents (215)	2.2	11	1	91 5	$\frac{428}{20}$	
Total vegetable food	6.1	34	6	345	1,610	
Total food	26.4	156	120	435	3,540	

The family here studied were in better circumstances than the majority of those described in this bulletin. The quantity of protein in the food per man per day was largely in excess of that called for by the commonly accepted dietary standard; the energy, however, was very close to that called for by the dietary standard for a man at moderate work. The excess of protein indicates that a considerable reduction might have been made in the amounts of such animal foods as meat, fish, and eggs, which furnished protein rather than energy. At the same time this would have reduced the cost of the ration, owing to the relatively large amount expended for animal foods as compared with vegetable and more particularly cereal foods. The 21 cents expended for tomatoes furnished but little actual nutriment, but doubtless added to the palatability of the diet.

## DIETARY STUDY OF A WASHERWOMAN'S FAMILY (NO. 194).

This family consisted of the mother, 55 years old, and two adult children, a son 21 years old and a daughter 36 years old. There were also two children of the latter, girls, one aged 11 and the other 6 years, and two children of another daughter who was out at service, namely, a girl 11 years old and a boy 4 years old. The weights of the members of the family were 135, 140, 160, 80, 42, 75, and 30 pounds, respectively. The father and the two sons-in-law were dissipated and did not live with the family. The persons included in the study were sober and economical, but in very poor circumstances; the children were in rags. All, however, were in good health. The mother and the daughter did washing to pay the rent, and the daughter earned \$2.50 a week in addition. The son was a truck driver and earned \$7 a week. Food was purchased for cash by the day at the small markets.

The study began April 14, 1897, and continued ten days. The number of meals taken was as follows:

	Means.
Man	30
Two women (60 meals $\times$ 0.8 meal of man), equivalent to	48
Two girls, 11 years old (60 meals × 0.6 meal of man), equivalent to	36
Girl, 6 years old (30 meals $\times$ 0.5 meal of man), equivalent to	15
Boy, 4 years old (30 meals $\times$ 0.4 meal of man), equivalent to	12
Total number of meals taken equivalent to	1.11
Total number of meals taken equivalent to	- 141
Equivalent to one man forty-seven days.	

Table 29.—Weights and cost of food and nutrients in dietary study No. 194.

Trinds amounts and next of feed for ten days		Cost, nutrients, and fuel value of food per maper day.				
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.	
ANIMAL FOOD.  Beef: Shank, fore, 1 pound, 8 cents (23); steak,	Cents.	Grams.	Grams.	Grams.	Calories.	
round, 7.75 pounds, 79 cents (29). Veal: Chops, 0.85 pound, 15 cents (54). Mutton: Side, 4.38 pounds, 56 cents (52). Vork: Fresh (as ham), 5 pounds, 33 cents (71); chops, 0.40 pound, 5 cents (60).		24		1	321 179	
Fish: Smelts, fresh, 1.50 pounds, 12 cents (110)	1.6	1 9	7 10 14	18	10: 9: 25:	
Total animal food	8.5	54	. 70	18	94	
VEGETABLE FOOD,	,					
Cereals: Barley, pearled, 0.50 pound, 2 cents (126); bread, 17.31 pounds, 51 cents (134); buns, 15.50 pounds, 75 cents (140); buns, hot cross, 4.32 pounds, 25 cents (141); cake, coffee, 1 pound, 10 cents (143); cake, mixed, 2.20 pounds, 10 cents (142); cake, fruit, 1.50 pounds, 15 cents (145)	4.0	34	19	223	1,23 43	
nips, 1.61 pounds, 3 cents (218); sauerkraut, 3 pounds, 8 cents (212)	. 1.3	7	1	48	23	
Total vegetable food	6.5	41	20	379	1,910	
Total food	15.0	95	90	397	2, 858	

The family here studied represents a type of those who need to live as economically as possible. The cost of the food, 15 cents per man per day, was reasonable, but the amounts of protein and energy were small and might have been increased with probable advantage. they been increased one-fifth by increasing the amounts of food materials purchased, but keeping the kinds and proportions the same, the cost of the daily ration would have been about 18 cents. The nutritive value of the ration could have been increased without raising the cost by substituting more economical materials for some of those purchased. One of the most expensive items of food used was yeal chops at 18 cents a pound. The most economical foods were as usual the cereals, although here there was considerable difference in the relative economy, the buns and cakes being much more costly sources of nutriment than the bread. A considerable variety of fresh vegetables was also used. If two-thirds the cost of these fresh vegetables had been expended for dried beans or peas, if the money expended for buns and cakes had been used to purchase bread at the price paid, and if the veal chops and mutton side had been omitted from the diet and the money spent for them had been expended for round steak at the prices paid, the quantity of protein and energy per man per day would have been increased 39 grams and 575 calories, respectively, without increasing the cost. If at the same time the woman had been able to do her marketing as skillfully as the woman in dietary studies Nos. 31, 155, and 180, still more nutriment would have been obtained for the same money. The changes suggested, it is believed, need not have made the diet less palatable or attractive. Had a still greater reduction in cost seemed necessary it might have been accomplished by diminishing still further the quantity of meat and increasing the amount of cereal foods correspondingly.

## DIETARY STUDY OF A STABLEMAN'S FAMILY (NO. 195).

This study was with a family consisting of the father, 44 years of age, the mother, 39 years of age; four daughters, aged respectively 21, 17, 11, and 9 years; and three sons, aged respectively 15, 13, and 4 years. The weights of the members of the family were 170, 135, 145, 155, 75, 60, 120, 100, and 25 pounds, respectively. The father earned \$7.50 a week truck driving, but spent a good deal of it for drink. The older daughter earned \$7 a week in a box factory, of which she paid \$4 to her mother; the second daughter was idle at the time of the study. The oldest boy was a plumber's apprentice, and from his wages paid his mother \$4 a week for board. Food was bought by the day on one week's credit at the smaller markets. The members of the family were in good health.

The study began April 16, 1897, and continued ten days. The number of meals taken was as follows:

	Meals.
Man	30
Two women (60 meals $\times$ 0.8 meal of man), equivalent to	48
Girl, 17 years old (30 meals × 0.7 meal of man), equivalent to	21
Boy, 15 years old (30 meals × 0.8 meal of man), equivalent to	24
Two children, boy 13 and girl 11 years old (60 meals × 0.6 meal	of
man), equivalent to	36
Girl, 9 years old (30 meals × 0.5 meal of man), equivalent to	15
Boy, 4 years old (30 meals × 0.4 meal of man), equivalent to	12
Total number of meals taken equivalent to	186
Equivalent to one man sixty-two days.	

Table 30.-- Weights and cost of food and nutrients in dietary study No. 195.

		Cost, nutrients, and fuel value of food per mar per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carboby- drates.	Fuel value.		
ANIMAL FOOD.  Beef: Round, 1 pound, 11 cents (29); shank, fore, 3 pounds, 15 cents (23); shank, hind, 5 pounds, 30 cents (24); steak, round, 5.50 pounds, 52 cents,	Cents.	Grams.	Grams.	Grams.	Calories.		
(29); corned, rib, 12.19 pounds, \$1.05 (8); roast, round, 6.81 pounds, 73 cents (22)  Pork: Bacon, 0.50 pound, 5 cents (59); ham, 5.25	4.6	43	63		763		
pounds, 75 cents (65); ribs, sparerib, 3.50 pounds, 25 cents (81). Fish: Cod, fresh, 1.50 pounds, 10 cents (87).	1.7	11 1	25		278 4		
Eggs, 7.44 pounds, 53 cents (117)  Butter, 1.38 pounds, 38 cents (118)  Milk, 20.12 pounds, 58 cents (124)	.9	5	5 9 6	7	71 84 105		
Total animal food	8.9	66	108	7	1,305		
VEGETABLE FOOD.							
Cereals: Rice, 0.50 pound, 4 cents (130); bread, 19.45 pounds, 81 cents (134); crackers, soda, 3.36 pounds, 14 cents (156); rolls, 1.25 pounds, 9 cents (164) Sugar, 7.25 pounds, 37 cents (169) Vegetables: Cabbage, 4.26 pounds, 5 cents (180); onions, 1 pound, 1 cent (195); potatoes, 21.19	1.7	17	4	102 53	525 217		
pounds, 34 cents (204); tomatoes, 8 pounds, 20 cents, (215); turnips, 4.25 pounds, 5 cents (218)	1.1	5	1	36	178		
Total vegetable food	3.4	22	5	191	920		
Total food	12.3	88	113	198	2, 225		

As in the preceding study, the quantities of protein and energy were less than are usually regarded as desirable for persons with moderate work; the cost also was quite small. However, the total cost of food during the study was \$7.60, or 10 cents more than the total income which the mother had for running expenses. As already indicated, the father drank so that the family received but a portion of his very limited wages, and food was purchased on one week's credit. The prices of the food materials were on the whole larger than were paid for similar materials by some of the other families studied who lived in the same region. The quantity of meats consumed was largely in excess of that used by many families in similar or even better circumstances. The money would have been more economically expended had one-half of that spent for meat been used to purchase cereals, and had the sum expended for the fresh vegetables been diminished one-half. With these changes the protein and energy per man per day would

have been increased by about 10 grams and 625 calories, respectively. If the diet had then been increased by about one-fourth, it is probable that the family would have been better nourished, although so far as could be seen their health at the time of the study was good. Of course, it must be borne in mind that the food consumption during so short a period does not necessarily give a fair indication of the normal food consumption of the family.

# DIETARY STUDY OF A TRUCKMAN'S FAMILY (NO. 196).

This study was carried on in a family consisting of the father, 52 years old; his wife, 46 years old; and two children, a boy of 15 and girl of 8 years, all Americans. Their weights were 125, 120, 75, and 50 pounds, respectively. The father, a truckman, had been in poor health and out of work for some time. The woman earned a little by washing, scrubbing, etc. They occupied two rooms, for which they paid \$7.50 a month rent. The family appeared poorly nourished. Food was bought in small quantities for cash and there was no visible waste.

The study began April 28, 1897, and continued ten days. The number of meals taken was as follows:

	Meals.
Man	. 30
Woman (26 meals × 0.8 meal of man), equivalent to	. 21
Boy, 15 years (30 meals $\times$ 0.8 meal of man), equivalent to	. 24
Girl, 8 years old (29 meals $\times$ 0.5 meal of man), equivalent to	. 14
Total number of meals taken equivalent to	_ 89
Equivalent to one man thirty days.	

Table 31.—Weights and cost of food and nutrients in dietary study No. 196.

Kinds, amounts, and cost of food for ten days.	Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.	
ANIMAL FOOD.  Beef: Tripe, 1.69 pounds, 12 cents (38). Mutton: Leg, 2.69 pounds, 34 cents (47) Pork: Loin, fresh, 2.13 pounds, 23 cents (61) Fish: Halibut, fresh, 1 pound, 22 cents (94) Eggs, 3.50 pounds, 30 cents (114) Butter, 1.06 pounds, 24 cents (118) Milk, 1.31 pounds, 3 cents (124) Milk, condensed, 3 pounds, 21 cents (125)	1.0 8	Grams.  11 4 2 7	Grams.  7 8 1 5 14 1 4	Grams,	Calories.  110 91 18 75 131 18 152	
Total animal foodvegetable food.	5.6	29	40	25	595	
Cereals: Flour, 0.50 pound, 3 cents (131); rice, 0.25 pound, 2 cents (130); bread, 5.82 pounds, 19 cents (134); crackers, fancy, 5 pounds, 50 cents (155); rolls, coffee, 2.25 pounds, 8 cents (143); rolls, sweet, 3 pounds, 22 cents (144)  Sugar, 8.76 pounds, 47 cents (169)  Vegetables: Beans, dried, 1.75 pounds, 10 cents (175); onions, 0.25 pound, 2 cents (195); peas, canned, 1 pound, 8 cents (199); potatoes, 4.11 pounds, 5 cents (204); turnips, 1 pound, 3 cents (218)	3.5 1.6	21	17	150 133	859 545	
Fruits: Raspberry jelly, 1 pound, 10 cents (231)	.3			9	36	
Total vegetable food	6.3	29	18	321	1,600	
Total food	11.9	58	58	346	2, 195	

This study is remarkable for the small amounts of protein and energy which, according to the statistics, were consumed per man per day. The protein was about half and the energy two-thirds the normal amount for a man at moderate work. The cost, 12 cents per man per day, would have been very moderate had the diet been sufficient, but to bring the diet up to the standard by use of the same materials in the same proportions would require an expenditure of from 18 to 24 cents. The food purchases of this family, as a rule, were not marked by wise economy. The meats purchased were the higher priced cuts and the prices per pound were considerably in excess of those paid by some other families for similar cuts. Another evidence of the injudicious selection of food when the income was so limited was the purchase of fancy crackers at 10 cents a pound and sweet rolls at 7.3 cents a pound instead of bread which might have been purchased at 4 cents, or stale bread at 2 cents a pound. The selection of vegetables also was not such as would give the largest amount of nutriment for the expenditure. Dried beans furnished the most nutriment for the money expended.

Had the family purchased less expensive cuts of meat, spent less for crackers, rolls, canned and green vegetables, and more for flour, rice, bread, dried beans, and potatoes, the quantities of protein and energy in the diet might have been increased without increasing the cost. If the same kinds of food as used had been increased in amount by about one-third to one-half, the family would doubtless have been better nourished and the cost would have been but 16 to 18 cents per man per day. All the family seemed poorly nourished and were rather sickly in appearance, hence anything which would increase the quantity of nutriment, even if it detracted to some extent from the variety of the diet, would have been advantageous. If, in addition to the changes already suggested, less had been expended for animal foods and more for cereals the diet would have been rendered still more nutritious although it might not have been quite as appetizing. Of course in cases like this it must be borne in mind that where the mother works out herself she has not the time and opportunity for the small domestic economies possible for a woman who remains at home. Meats, fancy breads and pastries, and a variety of vegetables are easily prepared in such a way as to be appetizing and palatable, while it requires some skill and thought to prepare the more common cereals so that they will be as attractive and appetizing.

### DIETARY STUDY OF A HUCKSTER'S FAMILY (NO. 197).

The family here studied consisted of the father, a native of Scotland, 50 years old; the mother, a native of Ireland, 35 years old, and six children, born in America—a girl 14, a boy 12, a boy 10, a girl 8, and a boy 2 years old, and an infant 3 months old. The weights of

the members of the family were 135, 140, 75, 65, 60, 50, 25, and 15 pounds, respectively. The father, a huckster, sold wild flowers and shrubs, making about \$4.50 a week. Neither the mother nor the children were able to add anything to the family income. The family occupied two rooms, for which they paid \$6.50 rent per month.

The study began April 28, 1897, and continued ten days. The number of meals taken was as follows:

M	eals.
Man	30
Woman (30 meals × 0.8 meal of man), equivalent to	24
Girl, 14 years old (30 meals $\times$ 0.7 meal of man), equivalent to	21
Two boys, 12 and 10 years old (60 meals × 0.6 meal of man), equiv-	
alent to	36
Girl, 8 years old (30 meals $\times$ 0.5 meal of man), equivalent to	15
Boy, 2 years old (30 meals $\times$ 0.4 meal of man), equivalent to	12
Infant, equivalent to	9
Total number of meals taken equivalent to.	147
Equivalent to one man forty-nine days.	

Table 32.—Weights and cost of food and nutrients in dietary study No. 197.

	Cost, nutrients, and fuel value of food per ma per day.				
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy-drates.	Fuel value.
ANIMAL FOOD.  Beef: Liver, 1.50 pounds, 10 cents (19); drippings,	Cents.	Grams.	Grams.	Grams.	Calories.
0.25 pound, 1 cent (13); corned beef, 2 pounds, 25 cents (3). Veal; Head-cheese, 1.50 pounds, 10 cents (67). Mutton: Chops, 2.75 pounds, 36 cents (45).  Pork: Bacon, 4 pounds, 48 cents (59); lard, 1 pound, 6 cents (69) Fish: Cod, fresh, 4 pounds, 24 cents (87); halibut,	1.7	13	20 32		240 310
3.06 pounds, 18 cents (93); chowder, clam, 6 pounds, 20 cents (86); mussels, pickled, 3 pounds, 15 cents (100).  Eggs, 4.84 pounds, 50 cents (114)  Butter, 1 pound, 29 cents (118)  Cheese, 0.50 pound, 6 cents (120)  Milk, 12.50 pounds, 33 cents (124)	1.0 .6 .1	12 6 1 4	2 5 8 2 4	5	88 71 75 23 78
Total animal food	6.8	39	73	11	885
VEGETABLE FOOD.  Cercals: Corn meal, 1.50 pounds, 3 cents, (151); oatmeal, 2 pounds, 11 cents (128); rice, 1 pound, 6 cents (130); bread, 28.46 pounds, \$1.02 (134); flour, prepared, 3 pounds, 12 cents (133); crullers, 7 pounds, 22 cents (167); macaroni, 1 pound, 8 cents (158).  Sugar, 5.37 pounds, 27 cents (169).  Vegetables: Greens, 4.13 pounds, 15 cents (190); lettuce, 1 pound, 10 cents (191); onions, 1.25 pounds, 5 cents (195); peas, split, 1.50 pounds, 9 cents (200); potatoes, 12.76 pounds, 17 cents (204); potatoes, 1 pound, 1 cent (205); scullions, 1 pound, 10 cents (213); tomatoes, canned, 2 pounds, 7 cents (216)	. 6	38	19	231 50 36	1, 279 205
Total vegetable food	5.4	45	20	317	1,670
Total food.	12.2	84	93	328	2, 555

This family might be classed among the very poor, the income being but 75 cents a day. The quantity of nutrients per man per day in the food which they consumed was scarcely what would be required, according to the usual standard, by a man at light work, and probably was not sufficient for the needs of the family, for although they appeared to be in good health they were not robust. Had the diet been increased about one-third it would doubtless have more nearly suited the requirements of the people nourished; the cost would then have been a trifle over 18 cents per man per day. The corned beef used was expensive for a family in such circumstances. Equally nutritious meat of similar character could have been purchased at half the price, as was evident from purchases made by other families studied who lived in the same region. The mutton was also perhaps more expensive than the family could afford. The 35 cents spent for clam chowder and pickled mussels would have furnished much more nutriment had it been expended for some of the cheaper cuts of beef. It is also doubtful if the use of so many eggs (31 dozen at 16 cents a dozen) was warranted, although they furnished as chean a source of nutriment as some of the meats used. A considerable variety of cereal products was used, the most expensive being the crullers at 7 cents a pound. The variety of green vegetables was, in this study as in the majority of those here reported, larger than the resources of the family would seem to warrant. Much the same suggestions for improvement as in the previous study could be made.

## DIETARY STUDY OF A LONGSHOREMAN'S FAMILY (NO. 198).

The Polish longshoreman's family here studied consisted of the father, 36 years old; the mother, 28 years old; and three children—a girl of 9, a boy of 5 years, and an infant 4 months old. The weights of the members of the family were 215, 165, 75, and 50 pounds, respectively, the weight of the infant not being stated. The father earned \$9 a week, and they paid \$10 per month rent for three rooms, two of which were light. Provisions were bought in small quantities for cash. The family appeared well nourished.

The study began May 8, 1897, and continued ten days. The number of meals taken was as follows:

	Meals,
Man	30
Woman (30 meals × 0.8 meal of man), equivalent to	24
Girl, 9 years old (30 meals $\times$ 0.5 meal of man), equivalent to	15
Boy, 5 years old (30 meals $\times$ 0.4 meal of man), equivalent to	12
Infant, equivalent to	9
Total number of meals equivalent to	90

Equivalent to one man thirty days.

Table 33.— Weights and cost of food and nutrients in dietary study No. 198.

Kinds, amounts, and cost of food for ten days.	Cost, nutrients, and fuel value of food per man per day.					
Kinds, another, and cost of foot for ten days.	Cost.	Protein.	Fat.	Carbohy-drates.	Fuel value.	
ANIMAL FOOD.  Beef: Steak, sirloin, 4.50 pounds, 54 cents (33); chuck, 4.8 pounds, 47 cents (27); round, 2 pounds, 30 cents	Cents.	Grams.	Grams,	Grams, 1	Calories.	
(29); soup piece, 2 pounds, 16 cents (23); corned, 3.50 pounds, 35 cents (3). Veal: Loin, 5 pounds, 42 cents (57).  Pork: Chops, 3.13 pounds, 33 cents (61); trimmings, 12 pounds, \$1.20 (82); bacon, 1 pound, 12 cents.	7.5	53	44	 	626	
(59); ham, smoked, 4.26 pounds, 48 cents (66) Fish: White, 1.50 pounds, 18 cents (113). Eggs, 3.44 pounds, 25 cents (117). Butter, 4 pounds, 80 cents (118).	2.7	26 2 6 1	160 1 5 52		1,594 17 71 487	
Cheese, 2 pounds, 24 cents (120).  Cream, 0.41 pound, 4 cents (123a).  Milk, 50,92 pounds, \$1.03 (124).  Milk, condensed, 1 pound, 7 cents (125).	3.4		10 1 31 1	39	130 9 555 46	
Total animal food	23.3	123	305	48	3, 535 —	
Cereals: Barley, 2 pounds, 10 cents (126); flour, 3.50 pounds, 10 cents (131); rice, 1 pound, 6 cents (130); bread, 42.32 pounds, \$1.78 (134); cake, 8.50 pounds, 95 cents (142).  Sugars, starches, etc.: Sugar, 12.87 pounds, 59 cents (169); olive oil, 0.44 pound, 4 cents (173).	10.0	77	21	496 195	2, 545 856	
Végetables: Cabbage, 7 pounds, 22 cents (180); horseradish, 0.50 pound, 7 cents (191); onions, 3.50 pounds, 15 cents (195); potatoes, 24.71 pounds, 37 cents (204). Fruits: Prunes, 4 pounds, 50 cents (237); raisins, 2	2.7	10	1	79	374	
pounds, 20 cents (238); jelly, 0.50 pound, 3 cents (231)		2	1	·		
Total vegetable food		= 89 212	29	840	4, 080 7, 615	

The results of this dietary study are among the most interesting of those here reported. The protein in the ration was nearly double and the energy more than double that of the commonly accepted dietary standard for men at moderate muscular work. It must be borne in mind, however, that the man and the woman were large persons and that the man was engaged at quite active work. Food was purchased not only in large quantity, but also in considerable variety, so that the cost per man per day was unusually high. If it had been desired, the cost could have been reduced in the same way as has been indicated in the discussion of previous studies. The family consumed a large amount of pork trimmings during the study, an average of 1.2 pounds a day, and an unusually large quantity of bread, averaging 44 pounds a day. One-third the protein and one-third the energy of the diet were obtained from these two articles, at a cost of about onefourth of the total. Even making allowance for the activity and size of the members of the family, it would seem that the diet was larger than was called for, and that a material reduction might have been made. It is difficult to understand how this food consumption, costing \$8.50 per week, could have been maintained for any length of time on the income of the father, which was but \$9 a week. It has been observed that among the families studied, and especially those of foreign birth, there was at times during dietary studies a tendency to change somewhat the ordinary mode of living. This change is sometimes one of increased and sometimes one of decreased food consumption. In the present case it seems hardly probable that the study represents the average normal food consumption of this family.

# DIETARY STUDY OF A CARPENTER'S FAMILY (NO. 199).

This study was made in a German family comprising the father, 48 years old, weighing 160 pounds; the mother, 48 years old, weighing 150 pounds; their daughter, 10 years old, weighing 75 pounds, and a boy boarder, 5 years old, weighing 40 pounds. All were in good health. The father, a carpenter, had steady work, but at rather low wages, earning \$9 per week. The mother earned \$1.50 per week washing. The young boy who lived in the family brought in \$1.25 a week for board. During half of the study he was away visiting his father. The family occupied three rooms, paying \$11 a month rent. Provisions were bought daily for cash at the small markets. The food was prepared in the manner to which they were accustomed in Germany.

The study began May 11, 1897, and continued ten days. The number of meals taken was as follows:

	Meals.
Man	21
Woman (30 meals × 0.8 meal of man), equivalent to	
Girl, 10 years old (30 meals $\times$ 0.6 meal of man), equivalent to	. 18
Boy, 5 years old (12 meals $\times$ 0.4 meal of man), equivalent to	. 5
Total number of meals taken equivalent to	. 68
Equivalent to one man twenty-three days.	

Table 34.—Weights and cost of food and nutrients in dietary study No. 199.

	Cost, n	lue of food	i per man		
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.  Beef: Chopped, 0.75 pound, 8 cents (29); shank pieces, 3.13 pounds, 25 cents (23); bologna, 1	Cents.	Grams.	Grams.	Grams.	Calories.
pound, 10 cents (1); tripe, 2 pounds, 12 cents (38); liver, 1.50 pounds, 12 cents (19). Pork: Shoulder, 3.50 pounds, 35 cents (77); salt, 1.14 pounds, 8 cents (73); shoulder, smoked, 1.50 pounds,	2, 9	25	12	1	220
15 cents (79); lard, 0.50 pound, 4 cents (69)	3.0	14	52		550
Fish: Flounders, 3 pounds, 15 cents (92). Eggs, 3.19 pounds, 27 cents (117). Butter, 0.25 pound, 7 cents (118).	1.1	3 7	6 4		10 85 35
Cheese, 1 pound, 10 cents (120). Milk, 10.32 pounds, 25 cents (124).	1.0	5 7	7 8	10	85 135
Milk, condensed, 1.75 pounds, 14 cents (125)	. 6	3	3	19	133 120
Total animal food	9,9	64	92	30	1,240

Table 34.— Weights and cost of food and nutrients in dietary study No. 199—Continued.

	Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protien.	Fat.	Carbohy-drates.	Fuel value.	
VEGETABLE FOOD.  Cereals: Oatmeal, 2 pounds, 10 cents (129); rice, 0.50 pound, 4 cents (130); flour, 2.86 pounds, 15 cents	Cents.	Grams.	Grams.	Grams.	Calories.	
(131); bread, 5.94 pounds, 17 cents (134); cake, coffee, 2 pounds, 10 cents (143); cracker dust, 0.50 pound, 5 cents (154); rolls, wheat, 1.37 pounds, 8 cents (164); pie, apple, 0.50 pound, 5 cents (160); pie, custard, 1 pound, 10 cents (161).  Sugar, 1.69 pounds, 10 cents (169).  Vegetables: Cucumbers, 1 pound, 9 cents (187); greens, 0.50 pound, 2 cents (188); onions, 1 pound, 5 cents (195); potatoes, 17.50 pounds, 19 cents (204); tomatoes, canned, 2 pounds, 8 cents (216); turnips, 1.19 pounds, 2 cents (219).	.4	32		196 33	1,045 135 325	
Total vegetable food	6, 0	41	12	299		
Total food	15.9	105	104	329	2,745	

The quantity of protein in the food consumed per man per day by this family was not far from the average found in the studies of farmers and mechanics in comfortable circumstances in different parts of the country. The amount of energy, however, was small. If the diet had been increased to some extent by the use of more oatmeal, rice, flour, and bread, and if some of the money expended for green vegetables and canned tomatoes had been used to buy more cereal foods, the diet could have been increased easily as regards both protein and energy with but little, if any, increase in the cost per man per day.

### DIETARY STUDY OF A PAINTER'S FAMILY (NO. 200).

This family consisted of the father, 23 years old; the mother, 20 years old; an infant, 5 months old, and 2 male relatives (boarders), one 28 and the other 23 years of age. The weights of the adult members of the family were 135, 89, 135, and 140 pounds, respectively. The father earned \$7 a week painting chairs in a factory. The two boarders together paid \$6 a week for board; one of them was idle during the time of the study. The family is representative of a class known locally as "furnished roomers." They paid \$2.25 per week for a single room 12 by 6 feet, with a bed, stove, table, and two chairs. There was no waste, all crumbs even being used up. Employment was unsteady, and the family were always in debt.

The study began May 22, 1897, and continued ten days. The number of meals taken was as follows:

M M	eals.
Three men	89
Woman (30 meals × 0.8 meal of man), equivalent to	24
Infant, equivalent to.	9
Total number of meals taken equivalent to	122
Equivalent to one man forty-one days.	

Table 35.—Weights and cost of food and nutrients in dietary study No. 200.

	Cost, nutrients, and fuel value of food per m per day.				
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.  Beef: Steak, chuck, 2 pounds, 20 cents (27); frankfurters, 2 pounds, 20 cents (14); meat, 1.50 pounds, 16 cents (20); steak, round, 2 pounds, 23 cents (29); skirting, 3 pounds, 15 cents (36); stew piece, 0.75	Cents.	Grams.	Grams.	Grams.	Calories.
pound, 6 cents (24). Mutton, 0.75 pound, 6 cents (44).  Pork: Chops, 5 pounds, 47 cents (61).  Eggs, 0.75 pound, 5 cents (117).  Butter, 2.75 pounds, 62 cents (118).	1.5	22 7 1	21 13 1 26	1	285 150 13 241 4
Milk, 1 pound, 2 cents (124) Milk, condensed, 2.50 pounds, 17 cents (125)	.1	3	2	15	92
Total animal food	5.8	33	63	16	785
VEGETABLE FOOD.				1	
Cereals: Bread, 20.50 pounds, 80 cents (134); bread, rye, 2 pounds, 10 cents (136); cake, coffee, 0.25 pound, 2 cents (143); rolls, plain, 1.75 pounds, 10 cents (164); rolls, sweet, 1 pound, 5 cents (144).  Sugar, 4.75 pounds, 28 cents (169)	2.6	26	5	152 52	776 214
cents (208); tomatoes, canned, 1 pound, 8 cents (216).	1.7	8	1	65	310
Total vegetable food	5.0	34	6	269	1,300
Total food	10.8	67	69	285	2,085

The quantities of protein and energy consumed per man per day by this family were but little more than half the amount ordinarily considered necessary for men at moderate work, and undoubtedly more food than this would be required to maintain the family in good physical condition. A certain grade of work might of course be done upon a small amount of protein and energy; but it has been observed that within certain limits the character of the work improves as the diet becomes more liberal.

The family was very poor, always in debt, and the utmost economy in expenditure of all kinds was necessary. But while the cost of the food per man per day was small, the expenditures were not in all cases the wisest, for a more nutritious diet could have been purchased for the same amount of money. The relative economy of the materials purchased during this study is illustrated by the figures in the following table, showing the quantities of nutrients and energy in 10 cents' worth of each at the prices paid per pound.

Table 36.—Cost of food materials per pound, and amounts of protein and energy obtained for 10 cents in each, in dietary study No. 200.

		In 1 pound.		In 1 pound. Amount bought for 10 cent		10 cents.	Total	
Kind of food material.	Price per pound.	Protein.	Fuel value.	Total food.	Protein.	Fuel value.	ex- pended during study.	
Beef:	Cents.	Pound.	Calories.	Pounds.	Pound.	Calories.	Cents.	
Chuck steak	10.0	0.166	735	1.00	0.17	725	20	
Frankfurters	10.0	. 196	1,170	1.00	, 20	1,165	20	
Round steak	11.5	. 190	895	. 87	. 17	775	23	
Skirting	5.0	. 161	1,040	2,00	. 32	2,075	15	
Stew piece	8,0	. 096	405	1.25	. 12	. 505	6	
Mutton, stew piece	8.0	. 135	1.445	1.25	. 17	1,805	6	
Pork, chops	9.4	. 134	1,270	1.06	. 14	1,355	47	
Butter	22.5	. 010	3,605	. 44		1,600	62	
Milk, condensed	6, 8	. 088	1,520	1, 47	. 13	2,235	17	
Bread:			,			,		
Wheat	3.9	. 092	1,215	. 2.56	. 23	3, 110	80	
Rye	5.0	. 090	1,180	2.00	. 18	2, 365	10	
Rolls	5.7	. 097	1,470	1.75	. 17	2,575	10	
Rolls, sweet	5.0	. 081	1,450	2.00	. 16	2,905	5	
Sugar	5.9		1,860	1.70		3, 155	28	
Corn, canned	6.7	. 028	455	1.50	. 04	680	20	
Potatoes	1.0	. 022	385	10,00	. 22	3,875	28	
Green vegetables: Onions, rad-						2,010		
ishes	7.0			1.43	. 02	210	14	
Tomatoes, canned	8, 0	. 012	105	1.25	. 01	130	8	

The most expensive materials purchased in any quantity were the butter, beefsteak, canned corn, and radishes. The variety and amount of animal food and of green vegetables might have been reduced and the money thus saved used to purchase dried legumes, oatmeal, rice, flour, and bread. Apparently this would not have materially reduced the palatability of the diet, yet the quantity of nutrients would have been increased without increasing the cost.

Had the family spent their money for food as wisely as those described in dietary studies Nos. 178 and 185, they would not have been so poorly nourished. The use of stale bread instead of rolls, and of oatmeal and dried beans instead of canned corn, would have resulted in a considerable increase of nutriment, but not of cost.

#### DIETARY STUDY OF AN EXPRESSMAN'S FAMILY (NO. 201).

This study was made in a family comprising the father, 30 years old, weighing 135 pounds; the mother, 25 years old, weighing 130 pounds; a girl, 11 years old, weighing 75 pounds; a girl, 9 years old, weighing 60 pounds; a boy, 7 years old, weighing 55 pounds; a girl, 5 years old, weighing 38 pounds, and a boy 3 years and an infant 10 months old, whose weights were not stated. The father, an expressman, who owned his horse and wagon, earned on an average \$10 a week, although his income was variable. The mother apparently understood nothing about the purchasing and preparation of food; the home was poorly managed, and the table unattractive. Each member of the family had a small insurance, which required about \$1 a month to maintain. They occupied three rooms, for which they paid \$12 per month rent.

The study began May 22, 1897, and continued ten days. The number of meals taken was as follows:

М	eals.
Man	30
Woman (30 meals × 0.8 meal of man), equivalent to	24
Girl, 11 years old (30 meals $\times$ 0.6 meal of man), equivalent to	18
Girl 9 and boy 7 years old (60 meals $\times$ 0.5 meal of man), equiva-	
lent to	30
Girl 5 and boy 3 years old (60 meals $\times$ 0.4 meal of man), equiva-	
lent to	24
Infant, 10 months old, equivalent to	. 9
Total number of meals taken equivalent to	135

Equivalent to one man forty-five days.

Table 37. - Weights and cost of food and nutrients in dietary study No. 201.

W. A. A. A. A. A. S. Card Constant damage	Cost, nutrients, and fuel value of food pe per day.				per man
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.  Beef: Steak, chuck, 11 pounds, \$1,11 (27); round, chopped, 7.99 pounds, \$4 cents (29); shin, 2 pounds,	Cents.	Grams,		Grams.	
12 cents (23); corned, 6 pounds, 35 cents (3) Pork: Loin, fresh, 4.50 pounds, 41 cents (68); bacon, 1 pound, 10 cents (59) Eggs, 2.43 pounds, 30 cents (114) Butter, 3 pounds, 66 cents (118) Milk, 3.76 pounds, 11 cents (124)	1.5	45 7 3 0 1	18 3 26 1	2	529 196 40 242 22
Milk, condensed, 6 pounds, 42 cents (125)  Total animal food	9,8	62	90	33	1,235
Cereals: Bread, 20.40 pounds, \$1 (134); buns, 1.25 pounds, 5 cents (140); cakes, 2 pounds, 20 cents (149); cakes, coffee, 1 pound, 10 cents (143); cakes, sweet, 2.50 pounds, 20 cents (142); crackers, 0.25 pound, 2 cents (152); crullers, 7.75 pounds, 65 cents (157); rolls, water, 4.75 pounds, 25 cents (166); pic, apple, 0.50 pound, 5 cents (160).  Sugar, 10.63 pounds, 53 cents (169).		33	27	222 107	1, 297 439
Vegetables: Cabbage, 3 pounds, 5 cents (189); onions, 1.50 pounds, 8 cents (195); potatoes, 7.62 pounds, 10 cents (204); rhubarb, 3 pounds, 5 cents (210); soup greens, 1 pound, 2 cents (188).  Fruits: Apples, 1 pound, 10 cents (221); strawberries, 2 pounds, 10 cents (239)	.7	3	1 0	19	100
Total vegetable food	7.9	36	28	350	1,845
Total food	17.7	98	118	385	3,080

As regards the amounts of nutrients eaten little need be said concerning the results of this study. The quantity of protein and energy was not greatly below that found in the diet of working people in different parts of the United States and the cost was not large. By more judicious use of some materials at the prices paid and the substitution of more economical materials for some of those used the quantity of nutrients might have been sufficiently increased to meet all demands of the body without increasing the cost. Care and skill in the preparation of the food could have made this diet much more attractive.

#### DIETARY STUDY OF A WAITER'S FAMILY (NO. 204).

This family comprised only the husband, 40 years old, and his wife, 35 years old, weighing 175 and 125 pounds, respectively. The husband was a waiter in a restaurant, and earned \$7 a week and board. He was at home for but three meals during the time of the study. The couple occupied one furnished room, for which they paid \$2.25 per week rent. Provisions were bought daily at the small markets. In the opinion of those making the study the couple belonged to the lowest type found in the city among those who claim to have a home.

The study began June 22, 1897, and continued ten days. The number of meals taken was as follows:

21.0	GSFIN.
Man	- 8
Woman (30 meals $\times$ 0.8 meal of man), equivalent to	24
Total number of meals taken equivalent to	27
Fauivalent to one man nine days	

Table 38.— Weights and cost of food and nutrients in dietary study No. 204.

Kinds, amounts, and cost of food for ten days,	Cost, nu	itrients, ar	nd fuel val per day.	ue of food	per man
Kinds, and the control to the contro	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.  Beef: Beef, 0.75 pound, 8 cents (20); steak, sirloin, 0.50 pound, 6 cents (33). Mutton, 0.75 pound, 7	Cents.	Grams.	Grams.	Grams.	Calories.
cents (43b)  Pork: Chops, 2.37 pounds, 30 cents (61); spareribs, 3  pounds, 18 cents (61); bacon, 0.50 pound, 6 cents	2, 3	15	20		247
(59) Eggs, 1.02 pounds, 10 cents (115) Butter, 0.43 pound, 9 cents (118) Milk, 7 pounds, 16 cents (124).	6.0 1.1 1.0 1.8	38 8 0 12	81 5 19 14	18	909 79 177 253
Total animal food	12.2	73	139	18	1,665
VEGETABLE FOOD,					
Cereals: Flour, 1 pound, 5 cents (131); bread, 1 pound, 5 cents (134); bread, dry, 0.25 pound, 1 cent (134); bread, rye, 1.50 pounds, 5 cents (137); biscuit, 0.25					
pound, 2 cents (139) Sugar, 2 pounds, 10 cents (169). Vegetables: Potatoes, 14.44 pounds, 31 cents (204);	2.0 1.1	20	3	115 101	581 414
tomatoes, canned, 2 pounds, 10 cents (204),	4.6	17	1	138	645
Total vegetable food	7.7	37	4	354	1,640
Beverages: Beer, 13.50 pounds, 45 cents (243)	5.0	3		78	335
Total food	24.9	113	143	450	3, 640

The quantities of protein and energy in this study approached quite nearly to the standard for persons at moderate work. The cost, however, 25 cents per man per day, might easily have been reduced by more careful selection of food, if such reduction had been desired. It should be mentioned, however, that the cost includes one item which doubtless should be included in many of the other studies, but for various reasons could not be ascertained, namely, the amount paid for beer. The family used, on an average, about a quart a day.

## DIETARY STUDY OF A LANDLORD'S FAMILY (NO. 205).

This family consisted of a man 55 years old, weighing 200 pounds, and his wife 50 years old, weighing 175 pounds. The couple lived very simply, occupying two rooms. They rented twenty-five rooms for which they paid \$80 per month, and sublet the rooms for about \$2 or \$3 each per week. Their income was probably not far from \$200 per month. They did all the work in the house themselves and were very thrifty, owning several pieces of property outside of the city. Provisions were bought for cash partly in quantity and partly by the day.

The study began June 1, 1897, and continued ten days. The number of meals taken was as follows:

	Meals.
Man	. 30
Woman (30 meals × 0.8 meal of man), equivalent to	. 24
Visitor	. 2
Total number of meals taken equivalent to	- 56
Equivalent to one man nineteen days.	

Table 39.—Weights and cost of food and nutrients in dietary study No. 205.

		Cost, nutrients, and fuel value of food per man per day.					
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy-drates.	Fuel value.		
ANYMAT YOUR							
ANIMAL, FOOD,	Cents.	Grams.	Grams.	Grams.	Calories.		
seef: Steak, 0.75 pound, 10 cents (33); round, 6 pounds, 87 cents (29); sirloin, 1 pound, 16 cents (33). ork; Ham. smoked, 4 pounds, 48 cents (66); head-	5.9	34			372		
cheese, 1 pound, 10 cents (67)	3.1	18	40		446		
Cggs, 3.94 pounds, 38 cents (115)	2.0	14	10		150		
Sutter, 1.75 pounds, 36 cents (118)	1.9	1	35		329		
Theese, 1 pound, 10 cents (120). Theese, cottage, 1.50 pounds, 5 cents (122).	.5	6 7	8 10	1 0	103		
filk, 19.69 pounds, 30 cents (124)	1.6	16	19	24	343		
lilk, condensed, 1.50 pounds, 11 cents (125)	, 6	3	3	19	118		
Total animal food	15.9	99	150	44	1,98		
THOUGH DIT DOWN							
VEGETABLE FOOD,							
ereals: Flour, 2 pounds, 10 cents (131); flour, prepared, 1.50 pounds, 5 cents (133); bread, 8.13 pounds, 42 cents (134); bread, rye, 1.50 pounds, 5 cents (136); cake, 0.88 pound, 10 cents (149); rolls, water, 0.25 pound, 1 cent (166); rolls, Vienna, 0.75 pound, 5 cents (165); pie, lemon, 0.50 pound, 10 cents (162)  ugars, starches, etc.: Cornstarch, 0.56 pound, 5 cents (172); sugar, 6.25 pounds, 27 cents (169); centes (172); sugar, 6.25 pounds, 27 cents (189); cucumbers, 0.50 pound, 2 cents (187); onions, 2.50 pounds, 9 cents (195); potatoes, 10.10 pounds, 29 cents (204); radishes, 0.50 pound, 1 cent (209); tomatoes, 1 pound, 5 cents (216); tomatoes, canned, 4 pounds, 13 cents (216); pickles, cucumber, 0.25 pound, 1 cent	4.6	34	7	215 161	1,086		
(202). ruits: Lemons, 0.25 pound, 2 cents (233); strawber-	3.3	8	1	57	27		
ries, 1.19 pounds, 9 cents (239)	. 6	0	0	2			
Total vegetable food	10.2	42	8	435	2,030		

The daily food consumption per man per day in this study was considerably in excess of the ordinary standard for a man at moderate muscular work. The weights of both man and woman were, however, above the average, and it is very likely that they required more than the average quantities of nutrients, judging by what has been found with other families in good health who performed similar amounts of work. It seems probable that the dietary could have been reduced about one-eighth and still have been sufficient for the needs of the consumers. Such suggestions for changes are based on theoretical considerations. The fact is recognized that individuals vary considerably in their requirements. However, the changes suggested in this and other studies seem warranted on the basis of average results. It is interesting to note that, in spite of the fact that the couple were quite well-to-do, the amount paid for food, 26 cents per man per day, was not excessive.

#### DIETARY STUDY OF A CARETAKER'S FAMILY (NO. 206).

This family consisted of the mother, born in Ireland, and three children born in America; the mother was 36 years old, and weighed 139 pounds; one daughter 19 years old weighed 110 pounds, and the other, 17 years old, weighed 100 pounds; the weight of the 4-year-old son was not reported. The mother earned \$16 a month cleaning offices. The elder daughter earned \$7 a week as telephone operator, and the younger daughter earned \$3 a week as book folder. The family paid \$9 per month for the rent of three rooms. During three days of the study they had a poor woman helping about the house, who took her meals with them. Provisions were bought by the day for cash at the small markets.

The study began June 15, 1897, and continued ten days. The number of meals taken was as follows:

Meals.
Three women (63 meals $\times$ 0.8 meal of man), equivalent to 51
Girl, 17 years old (30 meals $\times$ 0.7 meal of man), equivalent to 21
Boy, 4 years old (30 meals $\times$ 0.4 meal of man), equivalent to
Visitor
_
Total number of meals taken equivalent to
Equivalent to one man twenty-eight days.

Table 40.—Weights and cost of food and nutrients in dietary study No. 206.

Kinds, amounts, and cost of food for ten days.	Cost, nutrients, and fuel value of food per man per day.				
	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.			~		
Beef: Steak, round, 3.25 pounds, 48 cents (29); sir-	Cents.	Grams.	Grams.	Grams.	Calories.
loin, 1 pound, 15 cents (33); cottolene, 0.75 pound, 8 cents (12). 'ork: Sparerib, neck, 2.25 pounds, 8 cents (80); 'spareribs, 6.62 pounds, 47 cents (81); ham, boiled,	2.5	13	21		249
0.50 pound, 10 cents (64); ham, smoked, 3 pounds, 35 cents (66); frankfurters, 0.25 pound, 4 cents (14)	3.7	34	62	l	71€
ish: Cod, fresh, 0.75 pound, 7 cents (87)	.3	1 2	$\frac{0}{2}$		27
Sutter, 2 pounds, 44 cents (118)	1.6	0	28		260
4ilk, 22.78 pounds, 63 cents (124)         3uttermilk, 2.50 pounds, 4 cents (119)         4ilk, condensed, 1 pound, 7 cents (125)	2.3 .1 .2	12 1 2	15 0 1	18 2 9	268 19 54
Total animal food	11.1	65	129	29	1,585
VEGETABLE FOOD.					
ereals: Bread, 14.75 pounds, 75 cents (134); cakes, coffee, 1 pound, 10 cents (143); cakes, sweet, 0.50 pound, 5 cents (149); crackers, soda, 0.50 pound, 4	]				
cents (156); wheat, rolls, 4.50 pounds, 5 cents (167), ugars, starches: Cocoa, 0.50 pound, 22 cents (171);	3, 5	31	6	192	971
sugar, 8.50 pounds, 40 cents (169)	2.2	2	2	141	608
'egetables: Cabbage, 6.25 pounds, 15 cents (180); cauliflower, 2.94 pounds, 10 cents (183); cucumbers, 1 pound, 3 cents (186); onions, 2 pounds, 5 cents (195); peas, green, 0.75 pound, 5 cents (201);					
potatoes, 12.82 pounds, 38 cents (204). ruits: Apricots, fresh, 0.50 pound, 4 cents (224); cherries, 0.50 pound, 4 cents (227); gooseberries, 1.50 pounds, 8 cents (229); lemons, 0.75 pound, 5 cents (233); peaches, preserved, 1 pound, 10 cents (235); raspberries, jam, 1.50 pounds, 16 cents (240); straw-	2.7	8	1	50	247
berries, 2 pounds, 20 cents (239); watermelons, 0.50 pound, 3 cents (241)	2.5	1	1	24	112
Total vegetable food	10.9	42	10	407	1, 935
Total food.	22, 0	107	139	436	3,520

The quantities of protein and energy per man per day in this study are about equal to those found on the average in the dietary of farmers, mechanics, and other working people in comfortable circumstances in various parts of the country. The cost, 22 cents per man per day, was not excessive for the kinds and amounts of food purchased. If the family had so desired, the expense could have been reduced to some extent by a different selection of food materials.

### DIETARY STUDY OF A SAILOR'S FAMILY (NO. 209).

This study was made in a family comprising the father, 50 years old; the mother, 39 years old; three boys, one 17, one 5, and one 3 years of age, and two girls, one 15 and the other 10 years of age. The weights of the members of the family were 150, 200, 100, 35, 25, 88, and 50 pounds, respectively. All were in good health. The father worked on a tug boat, earning \$30 a month. The oldest son was a porter and paid \$4.50 a week to his mother for board. The family

occupied three rooms, for which they paid \$11.50 per month rent. Provisions were purchased daily at the small markets for cash.

The study began June 15, 1897, and continued ten days. The number of meals taken was as follows:

	Meals.
Man	. 23
Woman (30 meals × 0.8 meal of man), equivalent to	. 24
Boy, 17 years old (30 meals × 0.8 meal of man), equivalent to	24
Girl, 15 years old (30 meals × 0.7 meal of man), equivalent to	. 21
Girl, 10 years old (23 meals × 0.6 meal of man), equivalent to	. 14
Boy, 5 years old (23 meals $\times$ 0.4 meal of man), equivalent to	. 9
Boy, 3 years old (30 meals $\times$ 0.4 meal of man), equivalent to	. 12
Total number of meals taken equivalent to	. 127
Equivalent to one man forty-two days.	

Table 41.—Weights and cost of food and nutrients in dietary study No. 209.

		Cost, nutrients, and fuel value of food per man per day.				
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.	
ANIMAL FOOD.  Beef: Steak, skirt, 3.50 pounds, 24 cents (32); shin, 2 pounds, 10 cents (23); shank, 4.50 pounds, 25		Grams.		. Grams.	Calories.	
cents (24). Mutton, 9.75 pounds, \$1 (52). Pork: Chops, 5 pounds, 50 cents (61); spareribs, 3 pounds, 32 cents (61). Eggs, 3.70 pounds, 38 cents (115). Butter, 1.50 pounds, 30 cents (118). Cheese, 0.50 pound, 8 cents (121). Milk, 16.34 pounds, 39 cents (124). Milk, condensed, 1 pound, 7 cents (125).	3.8 2.0 .9 .7 .2 .9	32     12     6     0     1     6	21 4	9 6	244 62 130 23 126 38	
Total animal food	8.7	58	91	15	1,145	
Cereals: Barley, 0.44 pound, 2 cents (126); flour, 2.38 pounds, 7 cents (131); bread, 3.50 pounds, 22 cents (134); bread, stale, 11.25 pounds, 28 cents (138); crullers, 5 pounds, 35 cents (157); cakes, 3 pounds, 25 cents (149); doughnuts, 1.50 pounds, 10 cents (157); rolls, water, 2 pounds, 10 cents (166); pie, apple, 0.39 pound, 5 cents (160).  Sugars, starches, etc.: Molasses, 3.50 pounds, 8 cents (170); sugar, 3.25 pounds, 16 cents (169).  Vegetables: Beans, 2 pounds, 9 cents (176); corn, canned, 3 pounds, 12 cents (185); greens (soup), 2 pounds, 7 cents (188); onions, 1.25 pounds, 5 cents (195); potatoes, 15.88 pounds, 22 cents (204); peas, green, 0.75 pound, 10 cents (201); tomatoes, green, 0.75 pound, 10 cents (201); tomatoes,	3.4	29	21	179 62	1, 049 259	
canned, 4 pounds, 12 cents (216)	1.8	7	1	47	232	
Total vegetable food	5.8	37	-9-3	288	1,540	
Total food	14.5	95	113	303	2,685	

This family were apparently in good health and it may be that the food was sufficient. It is probable, however, that had there been a little more protein and considerable more energy the diet would have been more suited to their needs. With an increase of the same kinds of food materials the cost would necessarily have been greater, but had

the increased expenditure been for flour, bread, dried legumes, etc., the cost of the diet would not have been proportionately increased. Thus, if they had purchased 2 pounds more of beans and 22½ pounds of stale bread in addition to the other foods used, the diet would have furnished 122 grams of protein and 3,375 calories of energy per man per day, at a cost of 16 cents. Had they omitted the canned tomatoes, corn, soup greens, and green peas the expense would have been reduced 1 cent per man per day, with a reduction of only 2 grams of protein and 64 calories of energy. The cost of the diet was very moderate and in this, as in other cases, the family were justified in spending a sum reasonable in proportion to their income to render the diet attractive and palatable.

#### DIETARY STUDY OF A HOUSEKEEPER'S FAMILY (NO. 210).

This family consisted of the mother, 54 years of age; two adult sons, one 31 and the other 27 years old; two daughters, one 18 and the other 15 years old; and three grandchildren, a girl of 9, a boy of 6, and girl of 4 years. The mother, German born, acted as house-keeper or janitor for the building in which they lived, thus paying the rent of their flat, which was valued at \$11.50 per month. The sons were both at work, the younger earning \$12 a week. The older son paid \$3.50 a week for his board. The older of the two daughters earned \$5 a week in a flower store, and the younger \$2.50 a week book folding. Food was purchased daily for cash. The family were not in the best of health; none of the members seemed strong.

The study began June 20, 1897, and continued ten days. The number of meals taken was as follows:

M	eals.
Two men	51
Two women (60 meals $\times$ 0.8 meal of man), equivalent to	48
Girl, 15 years old (30 meals $\times$ 0.7 meal of man), equivalent to	21
Two children, girl 9 and boy 6 years old (51 meals × 0.5 meal of	
man), equivalent to	26
Child, 4 years old (30 meals × 0.4 meal of man), equivalent to	12
Total number of meals taken equivalent to	158

Equivalent to one man fifty-three days.

Table 42.—Weights and cost of food and nutrients in dietary study No. 210.

Kinds, amounts, and cost of food for ten days.		utrients, a	nd fuel va per day.	lue of food	per man
Kinds, amounts, and cost of food for ten days.	Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
ANIMAL FOOD.  Beef: Steak, chuck, 5.50 pounds, 51 cents (27); bologna. 1 pound, 5 cents (1). Mutton, 5.50 pounds.	Cents.	Grams.	Grams.	Grams.	Calories.
65 cents (52).  Pork: Head-cheese, 1 pound, 10 cents (67); loin, 7	2,3	17	20		256
pounds, 70 cents (61); bacon, 1 pound, 10 cents (59); ban, boiled, 0.50 pound, 10 cents (64).  Fish: Clam chowder, 2 pounds, 10 cents (86); salmon, 10 pound, 10 cents (104); weak (fresh), 6	1.9	12	21	i	273
pounds, 30 cents (112a); sturgeon, 0.50 pound, 10 cents (112) Butter, 1.75 pounds, 42 cents (118) Milk, 25,72 pounds, 60 cents (124)	1.1	7		1	52 121 158
Total animal food.	7,2		68	12	
VEGETABLE FOOD.	-	_	-		
Cereals: Bread, 15.25 pounds, 63 cents (134); bread, rye, 3.50 pounds, 15 cents (136); cake, 3 pounds, 30 cents (149); doughnuts, 1 pound, 5 cents (157); rolls, 1.50 pounds, 10 cents (166).  Sugar, 4 pounds, 21 cents (169)	2.3	18	6	113 34	592 139
2 cents (195); potatoes, 13.60 pounds, 30 cents (204); sauerkraut, 4.50 pounds, 20 cents (212); tomatoes, canned, 2 pounds, 8 cents (216) Fruits: Apple butter, 0.50 pound, 4 cents (223)	1.2 .1	4 0	1	26 3	132 12
Total vegetable food	4.0	13:3	7	176	875
Total food	11.2	65	75	188	1,735

This family was certainly insufficiently nourished. Like the family in dietary study No. 158 and that in No. 200, the diet furnished only little over half the normal amounts of protein and energy. It is not surprising, in view of this fact, that the family did not appear strong and robust. While there were no particular extravagances in the diet; the food materials could have been selected much more wisely. The suggestions which have been given in connection with some of the preceding studies apply equally well in this case. Under the conditions it appears that choice must be made between variety of food and quantity of nutrients, as the sum available for food was not great enough to secure both. The mother had comparatively little time to devote to the preparation of the food and to marketing, which doubtless rendered it more difficult in this case to secure the greatest possible returns for the money expended.

The relative economy of the different materials used is shown in the table following, giving the quantities of nutrients and energy in 10 cents' worth of each at the prices paid per pound.

Table 43.—Cost of food materials per pound, and amounts of protein and energy obtained for 10 cents in each, in dietary study No. 210.

	rice per ound.			Amounts bought for 10 cents.			Total amount ex-
	ound,	Protein.	Fuel value.	Total food.	Protein.	Fuel value.	pended during study.
Beef, chuck steak	Cents. 9.3	Pound.   0,166	Calories. 735	Pounds.	Pound. 0.18	Calories.	Cents.
Mutton	11.8	. 192	1,560	. 85	.14	1,315	65
Head-cheeseLoin	10.0 10.0	. 195 . 134	1,790 1,270	1.00 1.00	.20	1,790 1,275	10 70
Bacon	10.0	. 091	2, 795 1, 320	1.00	.09	2, 800 655	10
Fish: Clam chowder.	5, 0	,018	195	2,00	.04	380	10
Salmon Weakfish	10.0	.195	680 205	1.00	. 20 . 17	680 415	10
Sturgeon	20.0 24.0	.193	950 3,605	.50	. 09	475 1,505	10 42
MilkBread	2.3	.033	325 1, 215	4, 35	.14	1,385 2,935	60 63
Rye bread	4.3	. 090	1,180 1,675	2.33 1.00	. 21	2, 760 1, 675	15 80
Doughnuts	5. 0 6. 7	. 067	2,000 1,300	2.00 1.50	.13	4, 015 1, 950	5 10
Sugar Green vegetables: Beans,	5. 3		1,860	1.89		3,545	21
greens, and onions	3.5 2.2	. 022	385	2,86 4,55	.10	885 1,755	7 30
Sauerkraut	4.4	.017	125 105	2, 25 2, 50	. 04	280 260	20 8

Among the least economical foods in this study may be mentioned boiled ham at 20 cents a pound, clam chowder at 10 cents a quart, sturgeon at 20 cents a pound, cake at 10 cents a pound, sauerkraut at 10 cents a quart, and canned tomatoes at 8 cents a can. The most economical food was bread, but even this might have been purchased cheaper, judging by the facts brought out in other studies.

## SUMMARY AND DISCUSSION.

The financial circumstances of the families included in the dietary studies here reported varied widely. The regular income of one family was such that they might be called comfortably well-to-do; a few others had means at least sufficient for their actual needs, while there were some whose total income during the period of study was not equal to the cost of food alone. The large majority of them were in such circumstances that in all their purchases it was necessary that every cent should count. The results of the studies show a wide difference, however, in the ability of the families to make the most of the means at their disposal; some of them obtained ample nourishment at a reasonable cost, while others for the same or even a larger expenditure were not sufficiently nourished. A number of the poorer families were especially undernourished, but some of them obtained much more nutriment for the money expended than did others. It is interesting to compare the studies in these respects.

## PECUNIARY ECONOMY OF FOOD PURCHASED.

In the discussion of the studies in the preceding pages, some tables have been given showing the amounts of protein and energy obtained for 10 cents in the different food materials used. The studies for which such tables were provided are believed to be more or less typical, so that the remarks made concerning the economy of the purchases may be applied in a general way to all the studies. In order to show the variations in the prices paid by different families for similar food materials, and especially to illustrate the relative value and economy of different materials as sources of protein and energy, these tables are summarized here.

Table 44.—Cost per pound and protein and energy in 10 cents' worth of various food materials in some typical dietary studies.

The day to mind.	Dietary	Price paid per		unt for 10	cents,
Food materials.	study.	pand per pound.	Total weight.	Protein.	Energy
eef:		Cents.	Pounds.	Grams.	Calories
Chopped	186	10.0	1.00	86	89
Sirloin	154	17.6	. 57	42	55
Do	186 161	13.0	.77 1,25	59 97	76
Flank Fore shank	161	5, 6	1.79	104	97
Do	186	7,3	1.37	82	7.
Hind shank	178	4.6	2.17	94	87
Shank	172	- 4.0	2, 50	109	1,00
Neck	172	8.0	1.25	82	9,
Chuck steak	178 200	10.9	. 92	69	6
Do Do	210	10.0	1.00 1.08	75 81	7:
Round	200	11.5	.87	75	7
Do	186	12.0	. 83	73	7.
Skirting	200	5.0	2.00	146	2,0
Stew piece	200	8.0	1.25	55	5
Corned	178	5.2	1.92	159	2,6
Corned, canned	161 172	20.0	. 50 1. 25	64 117	5 7
Liver Tripe, pickled	178	5.0	2.00	106	5
Suet	186	5,0	2.00	41	7, 0
eal:	200		2.00	**	,,,,,
Chops	154	12. 2	.82	. 74	6'
Do	186	12.5	. 80	73	6
Cutlets	186	14.0	.71	64	4
amb chops	161	13.7	.73	62	1, 1
Chops	161	18, 5	.54	1 39	9
Leg	172	7.2	1.39	95	1, 2
Neck	172	4.5	2.22	123	2, 1
Stew piece	200	8.0	1.25	77	1,8
Side	210	11.8	. 85	62	1,3
ork: Chops	161	11.2	. 90	67	1 1
Do	172	10. 0	1.00	61	1, 4 1, 2
Do	186	9.6	1.04	77	1.6
Do	200	9.4	1.06	65	1.3
Loin	178	7.7	1.30	97	2,0
Do	210	10.0	1.00	61	1.2
Head-cheese	210	10.0	1.00	89	1.7
Trimmings	178 154	8.0	1.25 .83	28 54	3,5
Ham, smoked	186	20.0	. 50 . 50	32	1, 3
Do	210	20.0	.50	46	6
Bacon	172	12.0	. 83	35	2.3
Do	210	10.0	1.00	41	2,8
Salt	161	7.0	1.43	12	5, 2
Do	172	9.8	1.02	9	3,7
Feet	172	6.0	1.67	120	2,3
Lard	186 200	6.0	1,67	1	7,0
ausage, Frankfurthicken	154	11.9	.84	89 74	1,1
***************************************	186	16.0	. 63	54	6

Table 44.—Cost per pound and protein and energy in 10 cents' worth of various food materials in some typical dietary studies—Continued.

	Dietary	Price	Amount for 10 cents.		
Food materials.	study.	paid per pound.	Total weight.	Protein.	Energy
Pish:		Cents.	Pounds.	Grams.	Calories
Cod, salt	186	6.0	1.67	145	52
Do	178	8.0	1.25	108	46
Cod, fresh	154 172	14.8 6.2	1.61	50 81	21 46
Shad	186	10.0	1.00	. 86	75
Bluefish	172 178	6.0.	1.67	76	35
Herring	178	5.4	1.85	163	1, 21
Salmon	210 210	10.0	1.00 2.00	89 78	68
Weakfish	210	5.0 20.0	.50	43	4:
Sturgeon Herring, smoked	154	10.0	1.00	93	7
Salmon, canned	161	20.0	. 50	49	4.
Sardines Clam chowder.	161	6.0	1.67	179	1,5
Oysters	210 154	5.0 9.1	2.00 1.10	16 30	38
utter	154	17.1	.59	90	2, 1
Do	161	18.6	.54		1, 9
Do	172	21.9	. 46		1,6
Do	178	18.4	. 54		1, 9
<u>D</u> o	186	20.0	. 50		. 1,8
Do	200 210	22.5 24.0	.44		1, 6 1, 5
lilk	154	2.6	3.85	58	1, 2
Do	161	2.7	3.70	55	1,1
Do	172	2.3	4, 35	64	1,3
<u>D</u> o	178	3.2	3.13	47	1,0
Do	186	2.3	4.35	64	1,4
Do Iilk, condensed	210	2.3 7.3	4.35 1.37	64 55	1,3 2,0
Do	172	10.0	1.00	40	1,5
Do	200	6.8	1.47	59	2,2
Doheese	154	14.3	.70	82	1,3
Do	186	17.0	.59	68	1, 1
ggs	154	13. 1 10. 6	.76	51 59	5 6
ggs. Do Tour	186 172	2.4	4.17	211	6,8
100	178	2.8	3,57	185	5, 9
Bread	. 154	4.4	2.27	95	2,7
<u>Do</u>	161	2.6	3, 85	190	4,8
Do	172	4.6	2.17	91 233	2,6
Do	. 178 186	2.1	4.76	91	$\begin{bmatrix} 5, 9 \\ 2, 6 \end{bmatrix}$
Do	200	3.9	2.56	107	3,1
Do Brend, rye Do	210	4.1	2.44	101	2,9
Bread, rye	. 161	3.0	3.33	136	3, 9
Do	. 200	5.0	2.00 2.33	82 95	2,3
Do	210 161	4.3 3.3	3.00	127	2,7
trackers, soda	161	4.7	2.14	95	4,1
rackers, soda Rolls, water	. 172	4.7 5.3	1.89	76	2,4
Rolls	. 200	5.7	1.75	77	2, 5
Do	. 210	6.7	1.50	61	1,5
ake ake, mixed	. 154 161	9.5	1.05 2.00	30 58	1,7
ake	186	12.0	. 83	23	1,8
Do	. 210	10.0		28	1,6
Buns Joughnuts gie, apple Do	. 178	4.4	2, 27	84	3, 3
oughnuts	. 210	5.0	2.00	60	4,0
Te, apple	. 161 186	20.0	1.00	8 14	1,2
ugar	. 154	5.3		1.3	3,4
Do	161	5.5	1.82		3,8
Do	. 172	4.6 5.7	2.17 1.75		4,0
<u>Do</u>	. 178	5.7	1.75		3,1
Do	. 186	4.8			3,8
Dο	200 210	5.9 5.3	1.70		3, 1
Do Cornmeal	. 186	4.0	1.89 2.50	104	4, 1
\$1ce	154	8.0	1, 25	45	2,0
Do	. 186	6.2	1.61	59	2,6
)atmeal	. 161	2.3	4.35	320	8,
dacaroni	. 154	6.0		101	2,7
Vermicelli	. 186 186	6.2	1.61	100 54	1,8
Do. Vermicelli Green vegetables.	154	4.6		13	1 2
100	. 161		4.17	27	5
Do	. 172	.8	12,50	66 97	1,6
Do	. 178		6.25		

Table 44.—Cost per pound and protein and energy in 10 cents' worth of various food materials in some typical dietary studies—Continued.

	Dietary	Price	Amount for 10 cents.		
Food materials.	study.	paid per pound.	Total weight.	Protein.	Energy.
Green vegetables.  Do. Potatoes  Do. Do. Do. Do. Do. Do. Do. Do. Sourckraut  Tomatoes, canned Do. Corn, canned Beans, dried Beans, dried Cabbage Greens, dandelion Greens, - Onions Pickles Spinach Tomatoes Spinach Tomatoes Spinach Tomatoes Spinach Tomatoes Spinach Tomatoes Jelly Nuts	200 210 154 161 172 178 186 200 210 154 161 210 210 210 210 210 210 86 186 186 186 186 186 186 186 186	Cents. 7.0 3.5 3.5 3.1 1.3 1.0 2.2 9.3 5.3 4.4 3.9 8.0 4.0 6.7 6.7 5.0 10.0 8.0 3.8 10.7	Pounds. 1.43 2.86 3.13 5.00 9.10 7.69 10.00 4.55 5.1.08 2.25 2.55 1.25 2.50 2.70 2.00 6.67 1.18 1.82 2.20 2.00 1.00 1.05 1.25 2.63 1.93 1.00	Grams.  45 32 51 95 77 77 100 45 5 8 18 14 19 273 204 11 44 36 16 14 9 9 36 45	Calories. 210 8855 1, 220 8855 1, 277 3, 695 2, 960 3, 875 1, 755 280 265 1300 260 6880 4, 280 3, 210 8353 335 4000 4101 1101 1101 1101 1, 600 1, 600

The beef used by the different families included in this table varied from canned corned beef at 20 cents a pound in one study to beef shank at 4 cents in another, and the quantity of protein and energy obtained for 10 cents ranged from 159 grams and 2,650 calories, respectively, in corned beef (not canned) at 5.2 cents per pound to 42 grams of protein and 555 calories of energy in sirloin steak at 17.6 cents per pound. When the quantities of both protein and energy obtained are taken into account, this sirloin steak was perhaps the most expensive meat purchased.

A characteristic difference between beef and pork is well illustrated by the figures in the table, the pork, as a rule, furnishing considerably less protein and considerably more energy for 10 cents than beef.

The price paid for fish and shellfish ranged from 5 cents a pound for weakfish and clam chowder to 20 cents a pound for sturgeon and canned salmon. The fish most economically purchased was canned sardines at 5 cents a pound, furnishing 179 grams of protein and 1,580 calories of energy for 10 cents. The least economical purchase was clam chowder at 5 cents a pound (10 cents a quart), furnishing 16 grams of protein and 380 calories of energy for 10 cents.

The price paid for butter varied from 17.1 to 24 cents a pound, and the energy in 10 cents' worth from 2,105 calories at the lower price to 1,505 calories at the higher price.

Milk was purchased at prices for actual delivery ranging from 4.6

to 6.4 a quart, and constituted a fairly economical source of both protein and energy.

The price at which sugar was purchased varied from 4.6 to 5.9 cents per pound, with corresponding differences in the fuel value of 10 cents' worth, ranging from 4,020 to 3,155 calories.

The greater number of the families studied used little or no flour, but the majority of them purchased considerable quantities of bakers' goods. It is interesting to note in the studies summarized in the table the differences in the price paid for such articles. In 10 cents' worth of stale bread for 2.1 cents a pound one family obtained 233 grams of protein and 5.920 calories of energy, while another family paid 5 cents a pound for rye bread, and obtained for 10 cents only 82 grams of protein and 2,365 calories. The price paid for cake varied from 5 cents to 10 cents a pound, with corresponding differences in the quantities of protein and energy in 10 cents' worth. Buns at 4.4 cents a pound were fairly economical food, while pie at 20 cents a pound furnished comparatively little nutriment for the money.

The family which made this latter purchase, however, obtained many of their food materials at very reasonable rates, and the pie, of which only a small amount was purchased, was probably deemed a deserved luxury. This same family obtained protein and energy very economically in oatmeal at 2.3 cents a pound. It is interesting to note the difference between the nutrients in oatmeal at this price and in rice at 6 cents a pound, or in almost any of the other food materials included in the table. At such a price oatmeal undoubtedly constituted one of the most, if not the most, economical sources of nutriment noted in these dietary studies.

The amounts of protein and energy in green vegetables, such as onions, soup greens, green corn, and the like, are small. While vegetables are more or less of a necessity, in order to provide bulk, to supply the body with mineral salts, and to add to the palatability and attractiveness of the diet, these purposes could probably be served as well by a small as by a great variety. It is a question, therefore, whether it was wise under the circumstances to purchase green vegetables in such variety as was observed in some of the studies. The amount of money spent for soup greens by some families was out of all proportion to their food value. They contain practically no nutriment, and as flavoring materials they were rather expensive at the prices paid; that is to say, it is possible to season soups so that they are palatable with condiments, etc., which cost less. For instance, celery seed could probably be used at less cost than the fresh vegetable. The matter is important chiefly as an illustration of the fact that the practice which is easiest may not be the most economical. It requires more thought and more knowledge to use the less common kitchen condiments, which would in the end be cheaper, than to buy and use the soup greens.

Comparatively few of the families studied made use of such economical materials as the dried legumes in their diet. In the two instances included in the above table the price paid per pound by one family was 3.7 cents, while another paid 5 cents. Even at the latter price there was ten to fifteen times as much protein and energy obtained for 10 cents as there was in canned corn, canned tomatoes, or green vegetables that were used by so many of the families. The economy in the use of the dried legumes and the cereals has been repeatedly pointed out on preceding pages, especially as substitutes for the very uneconomical materials mentioned. It has also been suggested that they might very readily take the place of at least a part of the meat that is so generally considered a necessity by the laboring classes. They supply the same ingredient, protein, as the meat and at a much lower cost.

As will be seen from the table, potatoes were purchased at prices ranging from 1 cent to 3.2 cents per pound, with a corresponding range of 100 to 32 grams of protein and 3,875 to 1,220 calories of energy in the amounts obtained for 10 cents.

Canned tomatoes, which seems to have been a favorite food material with many of the families studied, constituted one of the most costly sources of both protein and energy. Under the circumstances, perhaps this food product should be regarded principally as an appetizer since undoubtedly it rendered the diet more palatable and acceptable and thus doubtless increased the consumption of bread or other food of less marked flavor. Under some circumstances, when used in this way, it perhaps need not be considered as an expensive dish. Its use by families so poor and so undernourished as some of those included in these studies certainly seems unwise, as it simply took the place of other materials very much more nutritious and not unpalatable which could have been purchased for the same sum.

# SUMMARY OF AMOUNTS OF NUTRIENTS AND ENERGY IN FOOD CONSUMED PER MAN PER DAY.

The relative economy of the different dietaries may be shown by a comparison of the cost and the quantities of nutrients and energy per man per day in each. Results of the studies reported in this bulletin are thus summarized in Table 45, which also include for purposes of comparison the results of some similar studies previously reported.<sup>a</sup> For convenience in comparison the results have been grouped according to the amounts expended for food, the basis for each group having been adopted arbitrarily, as follows:

Group A includes those in which the cost per man per day is less than 13 cents.

a U. S. Dept. Agr., Office of Experiment Stations Bul. 46.

Group B includes studies in which the cost per man per day ranged from 13 to 16 cents.

Group C includes studies in which the cost per man per day ranged from 17 to 20 cents.

Group D includes studies in which the cost per man per day ranged from 21 to 23 cents.

Group E includes studies in which the cost per man per day ranged from 25 to 28 cents.

Group F includes studies in which the cost per man per day was more than 28 cents.

The results of each group have then been averaged together, and the averages compared with one another. In the last group, however, the variation in cost and in amounts of nutrients and energy obtained is rather too wide to include the individual families in an average that could be taken as representative of any class.

Table 45.—Summary of results of dietary studies made in New York City.

Sewing woman's family	Diet- ary No.		Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
Group B.—Studies in which the cost per man per day ranged from 13 to 16 cents.	178 185 158 200 210 155 195 196	per day was less than 13 cents.  Sewing woman's family Longshoreman's family Laborer's family Plumber's family Painter's family Housekeeper's family Longshoreman's family Stableman's family Truckman's family	9 9 10 10 11 11 12 12 12	57 86 86 64 67 65 94 88 58	41 79 76 81 69 75 105 113 58	237 285 368 213 285 188 257 198 346	Calories, 1, 585 2, 255 2, 570 1, 890 2, 085 1, 735 2, 415 2, 225 2, 195 2, 555
Per day ranged from 13 to 16 cents.		Average of 10 studies	11	75	79	271	2, 151
167     Carpenter's family.     13     89     78     296       171     Tanner's family     13     113     108     356       177     Truckman's family     13     98     83     337       187     Watchman's family     14     94     116     391       170     Caretaker's family     14     94     116     391       172     Foundryman's family     14     102     128     462       34     Watchman's family     14     87     96     296       51     Shopkeeper's family     15     81     109     355       161     Washerwoman's family     15     101     145     446       183     Sail rigger's family     15     127     98     446       194     Washerwoman's family     15     95     90     397       209     Sailor's family     15     95     90     397       209     Sailor's family     16     72     98     314       110     Salesman's family     16     72     98     314       110     Salesman's family     16     79     125     347       112     Tin roofer's family     16     84     114		GROUP B.—Studies in which the cost per man per day ranged from 13 to 16 cents.					
194   Washerwoman's family   15   95   90   397	167 171 177 187 170 172 34 51 161	Carpenter's family Tanner's family Truckman's family Watchman's family. Caretaker's family. Foundryman's family Watchman's family Watchman's family. Shopkeeper's family. Washerwoman's family.	13 13 13 13 14 14 14 15 15	89 113 98 79 94 102 87 81 101	78 108 83 74 116 128 96 109 145	296 356 337 346 391 462 296 355 446	2, 255 2, 305 2, 925 2, 555 2, 430 3, 065 3, 505 2, 465 2, 800 3, 590 3, 590 3, 260
GROUP C.—Studies in which the cost per man per day ranged from 17 to 20 cents.  38 Sailors' boarding house. 17 96 129 181 162 Longshoreman's family 17 95 100 283 32 Jeweler's family 18 101 106 296	194 209 35 110 112 159 188	Washerwoman's family. Sailor's family Dyer's family Salesman's family Tin roofer's family Washerwoman's family Bookbinder's family.	15 16 16 16 16 16	95 95 72 79 84 119 85	90 113 98 125 114 91 88	397 303 314 347 227 463 309	2, 855 2, 685 2, 455 2, 910 2, 335 3, 230 2, 480 2, 745
162     Longshoreman's family     17     95     100     283       32     Jeweler's family     18     101     106     296		GROUP C.—Studies in which the cost per man per day ranged from 17 to 20 cents.	15	94		352	2,779
201     Expressman's family     18     98     118     385       192     Butcher's family     19     123     113     312       111     Tin roofer's family     20     99     125     327       180     Carpenter's family     20     121     138     442	162 32 52 201 192 111	Longshoreman's family Jeweler's family Housekeeper's family Expressman's family Butcher's family, Tin roofer's family. Carpenter's family.	17 18 18 18 19 20 20	95 101 93 98 .123 99 121	100 106 104 118 143 125 138	283 296 509 385 312 327 442	2, 335 2, 480 2, 610 3, 435 3, 080 3, 115 2, 910 3, 590
Average of 8 studies		Average of 8 studies	18	103	120	342	2,914

Table 45,—Summary of results of dietary studies made in New York City—Continued.

Diet- ary No.		Cost.	Protein.	Fat.	Carbohy- drates.	Fuel value.
	GROUP D.—Studies in which the cost per man per day ranged from 21 to 23 cents.					
		Cents.	Grams.	Grams.	Grams.	Calories.
166	Carpenter's family	21	126	135	452	3, 625
107	Truckman's family	22	136	135	595	4, 250
106	Printer's family	22	116	124		3, 120
160	Truckman's family	22	120	145	397	3,470
206	Caretaker's family	22	107	139		3,520
31	Carpenter's family	23	151	154	459	3, 935
47	Truckman's family	23	104	129	344	3,030
96	Laborer's family	23	139	119	345	3,090
108	Caretakers in day nursery	23	122	158	394	3, 585
	Average of 9 studies	22	125	138	420	3, 514
	GROUP E.—Studies in which the cost per man per day ranged from 25 to 28 cents.					
204	Waiter's family	25	113	143	450	3,640
33	Sailor's family	26	140	145	558	4, 190
186	Fruit vender's family	26	141	164	377	3, 650
193	Sail rigger's family	26	156	120	435	3,540
205	Landlord's family	26	141	158	479	4,010
168	Housekeeper's family	27	131	206	450	4, 295
97	Porter's family	28	142	142	444	3,720
31	Torter stammy		1.1~	174	333	0,720
	Average of 7 studies	26	138	154	456	3,864
	GROUP F.—Studies in which the cost per man per day was more than 28 cents.					
30	Mechanic's family	32	153	139	528	4,085
122	Mission worker's family	37	143	205	543	4, 725
154	Cable gripman's family	36	171	171	460	4, 175
147	Builder's family	41	187	219	723	5,770
198	Longshoreman's family	41	212	334	888	7,615
109	Builder's family	42	204	264	714	6, 220
109	Duract Stanning	45	204	20.3	11.1	0, 220
	Average of 6 studies	38	178	222	643	5, 432
			1			

As would be expected, the results show that the families expending the least for food received the least nourishment. Thus the average of Group A shows that among the families included 11 cents provided but 75 grams of protein and 2,150 calories of energy, while in Group F, at an average of 38 cents, there were obtained 178 grams of protein and 5,432 calories of energy. The difference in amounts obtained is not, however, proportionate to the difference in expense. Thus in the average of Group A each cent expended for food purchased about 6.8 grams of protein and 190 calories of energy, while in the average of Group F the amounts obtained for each cent were 4.7 grams of protein and 138 calories of energy, indicating that where there was less to spend there was greater economy in the purchase of food.

The figures in the table also illustrate what has already been pointed out regarding the differences in the ability of different families to provide for themselves economically. Thus in dietary study No. 48 there were obtained for 9 cents a day 57 grams of protein and 1,585 calories of energy, while for the same expenditure the family in dietary study No. 178 obtained 86 grams of protein and 2,255 calories of energy. On the other hand, practically the same amounts of nutrients and energy as in the latter case cost 12 cents for the family in dietary study No. 195. It is interesting to observe also that the family included in dietary

study No. 185 actually obtained for 10 cents more protein and energy than did the family in dietary study No. 188 for 16 cents. Other similar instances of differences in the economy of food purchases might be cited, but the above serve to indicate how one family may be well nourished while a neighboring family, expending for their food as much money per man per day, may be undernourished.

In Table 46 the results of the dietary studies in New York City here reported have been summarized by the averages of the various groups, Group A representing the smallest diet and Group F the most liberal diet observed. For the sake of comparison the results of studies with other persons or groups of persons under various conditions have also been included, as well as the commonly accepted dietary standards representing the average physiological demands of persons of different amounts of muscular work.

As already explained, the fuel values of these dietaries were calculated by use of the old factors, which allow 4.1 calories per gram of protein and carbohydrates and 9.3 calories per gram of fat. In the following table the results as thus calculated are summarized, and also the fuel values, as computed by use of the new factors previously mentioned, which are somewhat smaller, allowing 4 calories per gram of protein and carbohydrates and 8.9 calories per gram of fat. The quantities of digestible protein have also been computed and are given in comparison with the quantities of total protein in the various dietaries.

Table 46.—Comparison of the results of dietary studies in New York City with those of people in different conditions of life, and with dietary standards.

		Prot	ein.	Fuel	value.
	Cost.	Total.	Digest- ible.	By old factors.	By new factors.
	. Cents	Grams.	Grams.	Calories.	Calories.
Group A (smallest diet found in present investiga- tion), average of 10 studies	11	75	69	2, 151	2.087
Group B, average of 19 studies	15	91	86	2,779	2,692
Group C, average of 8 studies		103	95	2,944	2,848
Group D, average of 9 studies		125	115	3,514	3,408
Group E. average of 7 studies	26	138	127	3,864	3,747
Group F (most liberal diet found in present inves-					
tigation), average of 6 studies.	38	178	164	5, 432	5,260
Average of 17 mechanics' families	20	$\frac{106}{102}$	98 94	3, 454 3, 514	3, 343
Average of 11 farmers' families  Average of 18 professional men's families	26	102	99	3, 406	3, 407
Average of 4 poor families in Pittsburg	15	100	92	3, 261	3, 161
Average of 25 Bohemian families in Chicago	20	139	128	3, 483	3,376
Average of 5 Unorthodox Russian Jew families in				-,	
Chicago	22	144	132	3,044	2,954
Average of 11 Orthodox Russian Jew families in			440		
Chicago	19	122	112	3,041	2,951
Average of 4 Italian families in Chicago	17 22	100 108	9 <u>9</u>	3,008 3,170	2,917 3,067
Average of 5 French-Canadian families in Chicago Average of 25 families (previously studied) in desti-		103	55	5,170	5,007
tute circumstances in New York and elsewhere	9	84	77	2,653	2,573
(a) (a) (a) (a) (b) (b) (b) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c		0.		2,000	2,010
DIETARY STANDARDS.					
Man without muscular exercise (Atwater)		90	83	9 450	
Man with light to moderate muscular work (At-		50	00	2,400	
water)		112	103	3, 050	
Man with moderately active muscular work (At-					
water)		125	115		
Man with hard muscular work (Atwater)		150	138		
Man with very hard muscular work (Atwater)		175	161	5,500	

The results of the studies in New York City, when compared with those of similar studies and with the suggested dietary standards, indicate that a considerable number of the families were undernourished. Thus of the 59 families included in the summary, 29 in Groups A and B averaged searcely 90 grams of protein and 2,350 calories of energy per man per day, while the 8 families in Group C were also somewhat below the normal in the quantity of protein and considerably below it in the quantity of energy obtained. The 9 families in Group D obtained just about what is called for by the standard for a man at moderate work. The remaining 13 families in Groups E and F probably obtained more than they actually needed.

The families included in these studies in New York did not obtain as much nourishment for the money expended as was obtained by families in somewhat similar circumstances in other places. The 4 poor families in Pittsburg, included in the table, for 15 cents obtained 6 grams more protein and 500 calories more energy than were obtained for the same sum by the families included in Group B. A more striking contrast is found in the results of the studies among the families studied in Chicago, also included in the table.

An interesting comparison can be made between the results of the studies in New York and those made in Edinburgh, Scotland, and York, England, previously mentioned, among families in very much the same circumstances as those of the studies reported here. Averaging the 59 studies in New York City, the diet furnished about 110 grams of protein and 3,200 calories of energy per man per day, at a cost of 10 cents. The average of 16 studies of laborers' families in Edinburgh, carried on by Paton, Dunlop, and Inglis, shows about 100 grams of protein and 3,000 calories of energy per man per day, at a cost of 14 cents, and the average of 18 families in York, studied by B. S. Rowntree, shows 95 grams of protein and 2,900 calories of energy per man per day, at a cost of 14 cents.

#### CONCLUSION.

While the dietary studies of the poor in the congested districts of New York City are still too few in number to warrant sweeping conclusions, nevertheless they unmistakably indicate that a large portion of the laboring classes of those regions are undernourished. This condition, however, in the majority of cases was not due to a lack of means for obtaining sufficient nourishment; the difficulty was rather in the ignorance regarding the proper selection, purchase, and preparation of food materials. There are numerous illustrations in the studies of the fact that it was possible for even the poorer families to

<sup>&</sup>lt;sup>a</sup>A Study of the Diet of Laboring Classes in Edinburgh, pp. 44-56.

<sup>&</sup>lt;sup>b</sup>Poverty, a Study of Town Life, pp. 394-413.

obtain sufficient nourishment at a reasonable cost. There was searcely one case in which it was not easily possible, by a more judicious selection of food materials, to get more nutriment for the money expended than was obtained. Several instances have been pointed out in which some families were getting considerably more than others for the same expenditure.

In a number of cases the increase in nutritive value of the diet could have been obtained, perhaps, only by some sacrifice of variety, which might have made the diet less palatable. This, however, would depend largely upon the skill with which the more economical food materials were prepared for the table. While variety in the diet under some circumstances helps to increase the digestibility of food materials, still it is a question whether the variety found in some of the studies was of any special advantage in this respect. The extent to which variety must give place to actual nutritive value in the selection of foods is a question that must be settled by each family according to its circumstances.

Suggestions regarding the improvement of the food habits of the city's poor can be made here only in the general way in which they have been given in preceding pages. What was said in this connection in the report of the former studies in New York City" is directly applicable here.

From the results of all the studies, both those here given and those previously reported, it is quite evident that what is needed among these families more than anything else is instruction in the way to make the little they have go the farthest. This can best be done by concrete examples, by personal visitation and supervision of the purchase and preparation of food. In this there is a wide field for the operations of organizations such as the one which cooperated in making these studies, and a considerable amount of valuable work of this nature has already been undertaken.

Certain it is that improvements in the selection of food so as to secure more and better nutriment at less cost, in the cooking so as to make palatable dishes from inexpensive materials, and in the setting of the table so as to make it an attractive feature of home life, will be important means for the material and moral uplift of families like those whose dietary practice is described in this and the previous report.

a U. S. Dept. Agr., Office of Experiment Stations Bul. 46, pp. 63-65.

# APPENDIX.

As was explained on page 9, the percentages of nutrients assumed for the different food materials used in the dietary studies are given in the table following. These are all taken from a publication of this Office, giving average composition of American food materials, but are included here in order that the present bulletin may contain all the data used in the computations of the results here reported.

The numbers in parentheses given in connection with each food material in the detailed tables of the dietary studies on the preceding pages correspond to the numbers in the column headed "Ref. No." in the table below, and the values used for calculating the amounts of nutrients in any food material may be readily found.

Table 47.—Percentage composition of different food materials used in computing the nutrients in the food consumed in dietary studies in New York City.

Ref.   No.	Kind of food material.	Dietary studies in which used.	Protein.	Fat.	Carbohy-drates.
	Beef:	1	D	n	I D
7 1		151 150 100 010	Per cent.	Per cent.	Per cent.
1 0	Bologna	154, 159, 199, 210	18.2	19.7	
	Corned	170	14. 4	19. 4	
3	Do	171, 187, 188, 197, 198, 201	14.3	23. 5	
4 4a	Corned brisket	155, 158, 178	18.3	24.7	
	Corned flank	183	14.6	33.0	
5	Corned, canned	177	28.9	13. 7	
6	Do	161	28.2	15. 2	
7	Corned plate	160	11.7	35, 8	
8	Corned rib	195	13.7	41.9	
9	Do	177	17.5	26, 6	
10	Corned rump	168	14.3	22.0	
11	Corned shoulder	167	28.9	13.7	
12	Cottolene	180, 206		100, 0	
13	Drippings	197	4.1	82.1	
14	Frankfurters	200, 206	19.6	18, 6	1.1
15	Gelatin	154	91.4	. 1	
16	Heart	183	14.8	24.7	
17	Kidney	180	13.7	1.9	
18	Liver	159, 172	20.7	4, 5	1.5
19	Liver sausage	160, 166, 197, 199	20.7	4.5	1.5
20	Meat	200, 204	14.8	18.1	
21	Neck	172, 177, 183	14.5	11.9	
22	Roast	195	22.3	28, 6	
23	Shank, fore	155, 159, 160, 161, 166, 168, 177,	12.8	7.8	
		180, 186, 194, 195, 198, 199, 201,			
		209.			
24	Shank, hind	172, 178, 200, 209	9.6		
25	Do	195	20.9	11.5	
26	Shoulder	180, 185	16.4	9.8	
27	Steak, chuck	155, 158, 166, 168, 171, 177, 178,	16.6	10.1	
	,	198, 200, 201, 210.			
28	Do	183	18.5	18.0	
29	Steak, round	158, 159, 160, 162, 166, 167, 180,	19.0	12.8	
	,	186, 188, 189, 193, 194, 195, 198,	20.0	1210	
		199, 200, 201, 205, 206.			
30	Do	171	16.4	6.0	

a U. S. Dept. Agr., Office of Experiment Stations Bul. 28, revised.

Table 47.—Percentage composition of different food materials used in computing the nutrients in the food consumed in dietary studies in New York City—Continued.

					1
Ref. No.	Kind of fcod material.	Dietary studies in which used.	Protein.	Fat.	Carbohy- drates.
	Beef-Continued.		Per cent.	Per cent	Percent.
31	Steak, round, chopped	166, 167	20.3	13.6	1 Cr.ccnt.
32 33	Steak, skirt	209 154, 159, 162, 166, 168, 186, 187,	19.7	17. 7	
<i>త</i> ర :	Steak, sirloin	188, 192, 198, 204, 205, 206,	16.5	16.1	
34	Do	167	13.3	42.3	
35	Steak, skirt	159, 161, 177	17.0		
36 37	Steak, skirt Steak, skirting Suet	155 167 186	16.1 4.7	17. 5 81. 8	
38 29	Tripe	159, 161, 177. 185, 200. / 155, 167, 186. 155, 177, 178, 196, 199. 158, 177	11.7	1 ')	
	Dô	158, 171	16.8	8.5	
39a	Lamb:	197	7.5		
40	Breast	158	19.1	23, 6	
41	Chops Leg	161	18.7	28. 8	
42   43	Do.		15.9 19.2	13.6	
43a	DoShoulder	187	18.1	29.7	
4.53	Mutton:	204			
43b	Side	160 168 200 201	13. 0 13. 5	24, 0 28, 3	
45	Chops	160, 168, 200, 204 161, 177, 197	16.0	33.1	
46	170	168, 192 168, 171, 196	16.0	24, 1	
47 48	Leg	168, 171, 196 172	18.5 15.1	18.0	
49	Neck shoulder Do.	172	12.3	17, 9	
50	shoulder	158	17.7	19.9	
51 52	Side	158. 177, 192. 194, 209, 210.	13.7 16.2	15.5 29.8	
	Veal:		10. 2		
58	Breast	162	19.5	14.0	
51 55	Chops	162 154, 186, 194 186, 193	19.9 20.1	10.8	
100	Chops Cutlets Head-cheese ' Leg	197			
56	Leg	197 193	15.5	7.9	
57 58	Loin	198 170	16.6 15.1	9.0	
93	Pork:	1	10. 1	0.0	
59	Baeon	158, 159, 162, 168, 172, 183, 195, 197,	9.1	62.2	
60	Chops	198, 201, 204, 210. 154, 158, 161, 178, 180, 183, 186, 194, 159, 166, 171, 172, 196, 198, 200, 204,	16.6	30.1	
61	Do	159,166,171,172,196,198,200,204, 209,210.	13.4	24.2	
62	Feet	172	15.8	26.3	
63	Feet, pickled	183.	16.3	14.8	
64 65	Ham, boiled Ham, smoked Do.	183. 155, 177, 180, 206, 210. 168, 180, 195. 154, 162, 166, 171, 186, 187, 192, 198,	20.2 16.3	22. 4 38. 8	
66	Do	154, 162, 166, 171, 186, 187, 192, 198,	14.2	33.4	
		200, 206.	10.5		
67 68	Head-cheese Loin, fresh.	205, 210	19.5 13.2	33.8 26.0	
69	Lard	201 155, 158, 166, 177, 186, 197, 199	10, 4	100.0	
70	Lard. Pig's head. Pork as fresh ham. Salt.	155, 170	13.4	41.3	
71	Pork as fresh ham	161 167 170 179	15.3 1.9	28.9 86.2	
72 73 74	170	133	7.4	59.6	
74	Campaga		13.0	44. 2 32. 5	1.1
76 77	Sansage meat. Shoulder Shoulder, fresh	170	17.4	6.0	
77	Shoulder, fresh	199	12.0	29.8	
18	Shoulder, salt	199. 171, 185. 158, 167, 199.	15.9	32.5	
79	Shoulder, smoked	206	13. 0 17. 3	26.6 31.1	
81	Sparerip roasi	206. - 155, 160, 170, 195, 206. - 178, 198. - 154, 180, 180, 187, 193.	16.6	30.1	
82	Trimmings Poultry: Chickens	178, 198	5.0	65, 0 16, 3	
83	Fish:	104, 100, 100, 107, 190	19.3	10.3	
81	Bluefish, fresh	162, 172	10.0	. 6	5.2
85	Clams	158	10.6	1.1	5.2
86 87	Cod	158 197, 210 193, 195, 197, 206	8.4	.0	0.7
88	Cod, boneless Cod, fresh	171 155, 162, 171, 172, 188, 192	27.7	.3	
89	Cod, fresh	155, 162, 171, 172, 188, 192	11.1 16.5	.2	
90 91	Do	154, 183. 167, 168, 178, 186, 188.	19.0	.4	
92	Cod, salt	199	5.4	. 3	
93	Halibut	197	18.6 15.3	5, 2 4, 1	
94 95	Halibut, fresh Herring, fresh	196. 177.178.	19.5	7.1	
96	Herring, smoked Mackerel	104	20.5	8.5	
97	Mackerel	159	10.2	4.2	

a Composition assumed.

Table 47.—Percentage composition of different food materials used in computing the nutrients in the food consumed in dietury studies in New York City—Continued.

Ref.	Kind of food material.	Dietary studies in which used.	Protein.	Fat.	Carbohy- drates.
98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114	Fish—Continued.  Mackerel, fresh Mackerel, salt Mussels, pickled Oysters Perch Pike Salmon Salmon, canned Do. Sardines Shad Shed Shed Sturgeon Weakfish Whitefish	160, 170, 197, 188, 198, 199, 188, 199, 188, 199, 188, 160, 161, 187, 210, 161, 193, 186, 187, 210, 161, 193, 186, 187, 191, 162, 210, 210, 210, 210, 198, 159, 162, 166, 167, 168, 170, 177, 178, 185, 186, 187, 193, 194, 196, 197, 201, 210, 210, 210, 210, 210, 210, 210	13, 4	7, 5 12, 1 7, 5 12, 1 9, 5 1, 8 1, 6 14, 0 1, 1 3, 0 10, 5	1,0
115 116 117 118 119 120 121 122 123 124 124 125	Do D	154, 204, 205, 206, 209, 155, 161, 171, 180, 183, 188, 192, 195, 198, 199, 200 Used in all 206, 205, 198, 199, 205, 1209, 205, 162		9.3	.4
126 127 128 129 130	Cereals: Barley Farina Oatmeal Do Rice	159, 161, 177, 185, 197	\$.5 11.0 16.1 16.7 \$.0	1.1 1.4 7.2 7.3	77.8 76.3 67.5 66.2 79.0
131 132 133 134 135 136	Flour Flour, low grade Flour, prepared Bread Bread, brown Bread, rye	1 197, 198, 199, 159, 166, 167, 172, 177, 178, 180, 193, 196, 198, 199, 204, 205, 209, 171, 180, 197, 205, Used in all 180, 159, 161, 167, 170, 187, 188, 193,	11, 2 14, 0 10, 2 9, 2 5, 4 9, 0	1.0 1.9 1.2 1.3 1.8 .6	74.9 71.2 73.0 53.1 47.1 53.2
137 138 139 140 141 142	Biscuit, soda Buns Buns, sweet Cake	200, 203, 210. 204 155, 158, 159, 161, 178, 185, 209. 159, 160, 161, 165, 204. 160, 167, 178, 185, 188, 194, 201. 167, 194. 154, 158, 159, 161, 185, 186, 188, 194, 198, 201, 210. 160, 162, 193, 194, 196, 199, 200,	9.6 10.9 9.3 5.1 7.9 6.8	. 6 1.3 18, 7 6, 9 4. 8 9, 0	48, 9 53, 6 52, 6 54, 2 49, 7 63, 3
144 145 146 147 149 150 151 152 153 154 155 156 157 159 160 161 162	Cake, jelly Currant loaf Cake, sweet Cakes, sweet Corn cake Corn meal Crackers Do Cracker dust Crackers, faney Crackers, soda Crullers Macaroni Muflins Pie, apple	194   168   168   168   201, 205, 206, 209   168   186, 197   189   199   199   196   161, 177, 185, 187, 195, 206   161, 177, 185, 187, 201, 209, 210   154, 150, 170, 186, 197   188	6. 8 6. 7 5. 9 6. 3 7. 9 9. 2 10. 7 10. 9 7. 10. 9 8. 7 10. 7 10. 7 10. 7 10. 7 10. 7 10. 7 10. 7 10. 7 10. 7	6,9 10,9 10,7 7,6 9,0 9,0 4,7 1,9 8,8 12,1 6,0 13,5 12,1,0 9,1 121,0 9,6 6,3 10,1 9,6	51, 2 61, 1 65, 9 57, 6 61, 8 61, 8 75, 4 71, 9 68, 7 77, 9 68, 7 77, 1 1 41, 8 42, 8 26, 1 51, 4

Table 47.—Percentage composition of different food materials used in computing the nutrients in the food consumed in dietary studies in New York City—Continued.

	,				
Ref. No.	Kind of food material.	Dietary studies in which used.	Protein.	Fat.	Carbohy- drates.
-	0 1 0 1 1		-		
164	Cereals—Continued. Rolls, plain	177, 180, 188, 193, 195, 199, 200	Per cent. 9.7	Per cent.	Per cent. 59.9
165	Rolls, Vienna		8. 5	2.2	56, 5
166	Rolls, water	167, 168, 170, 172, 201, 205, 209,	9.0	3.0	54.2
4.05	7) - 11 4	210.	0.4		F (1
167 168	Rolls, wheat	206 186	9.4 10.9	2.0	59. 4 72. 0
1000	Vermicelli Sugars, starches, and oils:	AWRESTANCE	10.0	0	70
169	Sugar	t sed in all			100.0
170	Molasses		2.4 21.6	28.9	69.3 37.7
171 172	Coeoa Cornstareh	168, 186, 192, 206	21.0	۵۰. ت	90.0
173	Olive oil	186, 198		100.0	
77-4	Vegetables: Asparagus	154	1 0	0	4) 4)
174 175	Beans	166, 170, 178, 180, 183, 186, 187,	$\frac{1.8}{22.5}$	1.8	3.3 59.6
110		196.			00.0
176	Do	209, 210	4.7	.3	14.6
177 178	Beans, string Do	155	2.3 2.1	.3	7.4 6.9
179	Cabbage	155, 158, 160, 161, 162, 166, 167, 168, 170, 172, 177, 178, 180, 187, 183, 185, 186, 192, 194, 195, 198,	1.6	. 3	5. 6
		168, 170, 172, 177, 178, 180, 187.		_	
180	Do	183, 185, 186, 192, 194, 195, 198, 201, 205, 206.	1.4	. 2	4.8
181	Cabbage sprouts	187	4.7	1.1	4.3
182	Carrots	187. 178, 187, 194.	1.1	.4	9.3
183 184	Cauliflower	206 158, 166, 167, 170	$\frac{1.6}{3.1}$	1.1	4.2 19.7
185	Corn. Canned.	159, 160, 161, 168, 188, 200, 209,	2.8	1.2	19.0
186	Cucumbers	155. 206	.8	. 2	3.1
157	Greens	199, 205. 167, 179, 186, 201, 209, 210. 166, 171, 177, 180, 183, 186, 187,	2.4	1.0	2.6
188 189	Do	166 171 177 180 183 186 187	4.2	. 6	10.6
		194.			
190	Do Horse-radish	197	1.8 1.6	.4	1.7
191 192	Leeks.	198 193	1.0	.4	5.0
193	Lettuce	154	1.2	.3	2. 9 2. 5
194	Do	197	1.0	.2	2.5
195	Onions	154, 155, 158, 159, 160, 161, 166, 167, 170, 171, 172, 186, 193, 194,	1.4	. 3	8.9
		195, 196, 198, 199, 200, 201, 205,			
102	Do	206, 209, 210.	1.6		0.0
196 197	Parsley	177, 178, 180, 183, 187 186, 192, 199	2.4	1.0	9, 9 10, 6
198	Parsnips	193	1.3	.4	10.8
199	Peas, canned	160, 180, 196. 160, 178, 187, 197.	3, 6	. 2	9.8 62.0
200 201	Peas, dried Peas, green	206, 209	24, 6 7, 0	1.0	16.9
202	Pickles, cucumber	194. 205	.5	. 3	2.7
200	Pickles, mixed	186	1.1	. 4	4.0
204 205	Potatoes Potatoes, cooked	Used in all	2.2	. 1	15.4 20.9
206	Potatoes, sweet	170	1.4.	. 6	21.9
207	Do	193	1.8	. 7	27. 4
208	Radishes	154, 200	1.3	. 1	5, % 1, 0
209 210	DoRhubarb	159, 205. 201.	.9 .	. 4	2.2
211	Salad	166	1.2		2.9
212	Sauerkraut	166, 180, 194, 210	1.7	.5	3. S 9. 2
213 214	Scallions Spinach	197	1.4 2.1		5.2
215	Tomatoes	158, 159, 161, 162, 166, 167, 168,	.9	. 4	3, 9
		170, 171, 172, 180, 186, 187, 192,			
216	Tomatoes, canned	193, 195, 205. 154, 158, 159, 160, 162, 177, 183,	1.2	. 2	1.0
	,	188, 197, 199, 200, 204, 205, 209,			
217	The second second second	210. 168, 187.	1.5	. 2	12.3
218	Tomato catsup Turnips	162, 170, 171, 178, 183, 185, 192,	1.3	. 2	8.1
		194, 195, 196.		4	5.7
219	Fruits:	172, 199	. 9	.1	0.7
220	Apples, dried	177	1.6	2, 2	66.1
221	Apple	159, 170, 193, 194, 201	. 3	. 3	10.8
223	Apple butter	166, 168, 180 210	.4 1.2	1	14. 2 58. 5
2021	Apricots	206	1.0		12.6
5522	Bânanas. Do.	206 154, 167	1.3	. 6	21.0 14.3
226	Cherries	161. 154, 206	1.0	.4	16.7
228	Currants	159, 161	1.5		12.8
229	Gooseberties	206	.4	. 6	9. 9

Table 47.—Percentage composition of different food materials used in computing the nutrients in the food consumed in dietary studies in New York City—Continued.

Ref. No.	Kind of food material.	Dietary studies in which used.	Protein.	Fat.	Carbohy- drates.
230   231   232   233   234   235   236   237   238   239   240   241   242   243	Fruits—Continued. Grapes Jelly. Jelly, currant Lemons Muskmelon Peaches, preserved Pears Prunes Raisins Strawberries Raspberry jam Watermelon Nuts Beer	196, 198	1.0 1.2 .2	Per cent. 1.2 7.1 .5 8.0 .6 .6 .1 31.5	Per cent. 14.4 59.8 67.5 5.9 9.3 10.8 12.7 73.3 68.5 7.0 59.8 2.7 6.7 11.5













